

**MARIJUANA COCAINE
PRESCRIPTION DRUGS**

**THE CONSUMPTION AND CONSEQUENCES
OF ALCOHOL, TOBACCO, AND DRUGS IN INDIANA:
A STATE EPIDEMIOLOGICAL PROFILE
2018**

INDIANA STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP



**RICHARD M. FAIRBANKS
SCHOOL OF PUBLIC HEALTH**

INDIANA UNIVERSITY
Center for Health Policy
IUPUI



TOBACCO

HEROIN

METHAMPHETAMINE

ALCOHOL

THE CONSUMPTION AND CONSEQUENCES OF ALCOHOL, TOBACCO, AND DRUGS IN INDIANA: A STATE EPIDEMIOLOGICAL PROFILE 2018

Developed by the Indiana State Epidemiological Outcomes
Workgroup, 2018

Our Vision

*Healthy, safe, and drug-free environments
that nurture and assist all Indiana citizens to thrive.*

Our Mission

*To reduce substance use and abuse
across the lifespan of Indiana citizens.*

Published by the Center for Health Policy,
Indiana University Richard M. Fairbanks School of Public Health,
Indiana University-Purdue University Indianapolis (IUPUI)

This document, written for state policymakers and community leaders, presents data and analyses to support the development of a framework for advancing the mission of the Indiana Substance Abuse Prevention System.

This document and the efforts described herein were funded by the Indiana Family and Social Services Administration/Division of Mental Health and Addiction through the Substance Abuse Prevention and Treatment Block Grant CFDA 93.959 from the Substance Abuse and Mental Health Services Administration.

For additional copies of this document, contact:

Center for Health Policy

IU Richard M. Fairbanks School of Public Health
Indiana University-Purdue University Indianapolis (IUPUI)
1050 Wishard Blvd.
Indianapolis, IN 46202
Phone: 317-274-2000
IUCHP@iupui.edu



This document is available via the World Wide Web and can be accessed and downloaded from the Center for Health Policy Web site (<https://fsph.iupui.edu/research-centers/centers/health-policy/epi-reports.html>).

BOOK TEAM

Marion S. Greene, PhD, MPH
Casey Balio, BA
Harold E. Kooreman, MA, MSW
Corey Jacinto, MPH
Katy Hilts, MPH

Cover Design and Layout: Susan Hill

STATEMENT OF REPRODUCIBILITY

Permission is granted, free of charge, to photocopy pages of this document that are not copyrighted. Permission to reproduce from government sources is traditionally freely granted by the U.S. Government. If the analysis included in this report is quoted, the source should be credited.

Printed in the United States of America by the Center for Health Policy at IUPUI, Indianapolis, Indiana

INDIANA STATE EPIDEMIOLOGY AND OUTCOMES WORKGROUP (SEOW)

Marion Greene, Chair

Assistant Professor
Department of Health Policy & Management
IU Richard M. Fairbanks School of Public Health

David Bozell

Assistant Deputy Director
Indiana Division of Mental Health & Addiction

Grace Dickinson

Substance Abuse Program Manager
Indiana Criminal Justice Institute

Bethany Ecklor

Youth Services Bureau Chief
Indiana Division of Mental Health & Addiction

Therese Harper

Primary Care Behavioral Health Integration Evaluator
Indiana Division of Mental Health & Addiction

Raven Helmick

Prescription Drug Overdose Epidemiologist
Indiana State Department of Health

Davetta Henderson

Assistant Deputy Director
Indiana Division of Mental Health & Addiction

Aubrey Kearney

Research Analyst
Indiana Department of Child Services

Harold Kooreman

Policy Analyst, Center for Health Policy
IU Richard M. Fairbanks School of Public Health

Vera Mangrum

Partnerships for Success Coordinator
Indiana Division of Mental Health & Addiction

Don McCay

First Sergeant, North Operations Commander
Drug Enforcement Section
Indiana State Police

Hope McMickle

Evaluation Specialist
Indiana Prevention Resource Center

Jason Murrey

Statewide Suicide Prevention Coordinator
Indiana Division of Mental Health & Addiction

Ari Nassiri

Data Analyst
Indiana Division of Mental Health & Addiction

Pamela Pontones

Deputy Health Commissioner and State Epidemiologist
Indiana State Department of Health

Michael Ross

Division Director of Substance Abuse Services
Indiana Criminal Justice Institute

Katelin Rupp

Director, Program Evaluation
Tobacco Prevention & Cessation Commission
Indiana State Department of Health

Christian Shue

Colonel
Indiana Air National Guard
Chief of Aerospace Medicine, 122d Medical Group

Mike Toles

Sergeant
Methamphetamine Suppression Section
Indiana State Police

Kelly Welker

Assistant Deputy Director of Addiction
Indiana Division of Mental Health and Addiction

Jeff Wittman

School Social Work & Foster Youth Specialist
Indiana Department of Education

SEOW SUPPORT TEAM

Casey Balio

Research Assistant, Center for Health Policy
IU Richard M. Fairbanks School of Public Health

Samantha Childress

Research Facilitator, Social & Behavioral Sciences
IU Richard M. Fairbanks School of Public Health

Jyosna Gutta

Center for Health Policy Intern
IU Richard M. Fairbanks School of Public Health

Harold Kooreman

Policy Analyst, Center for Health Policy
IU Richard M. Fairbanks School of Public Health



Improving Community Health Through Policy Research

About the Center for Health Policy

The Center for Health Policy (CHP) is the research hub of the Department of Health Policy and Management. Our mission is to generate evidence that informs decision-making in Indiana and beyond. CHP Fellows and staff conduct rigorous research and evaluation on health system performance and health policy issues, with a specific focus on: population health and analytics; substance misuse and mental health services; and public health systems and services research.

The CHP has a vibrant research portfolio including funding from the National Institutes of Health (NIH), the Agency for Healthcare Research and Quality (AHRQ), the Robert Wood Johnson Foundation, various state agencies in Indiana, and numerous other government agencies nationwide.

The Center is directed by Dr. Joshua Vest.

TABLE OF CONTENTS

Introduction.....	1
1. Executive Summary.....	3
2. Alcohol Use in Indiana: Consumption Patterns and Consequences	7
3. Tobacco Use in Indiana: Consumption Patterns and Consequences.....	27
4. Marijuana Use in Indiana: Consumption Patterns and Consequences	51
5. Opioid Use in Indiana: Consumption Patterns and Consequences.....	61
6. Stimulant Use in Indiana: Consumption Patterns and Consequences	79
7. Mental Health and Suicide in Indiana	95
8. Methods and Data Sources.....	107
APPENDIX I. Data Sources Recommended by the State Epidemiological Outcomes Workgroup (SEOW)	115
APPENDIX II. Substance Use Indicators At-A-Glance.....	117
APPENDIX III. Cluster Analysis.....	120

INTRODUCTION

In July 2005, Indiana's Office of the Governor received a grant from the U.S. Department of Health and Human Services' Center for Substance Abuse Prevention (CSAP) as part of CSAP's Strategic Prevention Framework State Incentive Grant (SPF SIG) program. The SPF SIG program represented a continuation of ongoing CSAP initiatives encouraging states to engage in data-based decision-making in the area of substance use prevention planning and grant-making.

This grant was made on the heels of an earlier CSAP State Incentive Grant (SIG), which laid much of the groundwork for this new initiative. A great deal of work was completed under the first SIG to assess substance abuse prevention services and develop a strategic framework to guide policymaking in this area for the 21st century. The final report summarizing the outcomes of this work, entitled *Imagine Indiana Together: The Framework to Advance the Indiana Substance Abuse Prevention System*, was prepared by the Governor's Advisory Panel within the Division of Mental Health and Addiction (DMHA), Indiana Family and Social Services Administration. This report is available from DMHA and the Indiana Prevention Resource Center at Indiana University Bloomington.

As a requirement of the SPF SIG initiative, the State established a State Epidemiological Outcomes Workgroup (SEOW) to facilitate data-based decision-making regarding substance use prevention programming through the collection, analysis, and reporting of available epidemiological data. After the end of the Indiana SPF SIG in 2010, the State decided to continue supporting the work of the SEOW as part of its long-term efforts to improve substance abuse prevention policy.

This report represents the 13th official *State Epidemiological Profile* completed by the SEOW. As in past years, we have updated the core set of analyses to reflect the most recent data available. In order to make the report most useful for state and local policymakers and service providers, we present detailed information and descriptive analyses regarding the patterns and consequences of substance use both for the state and, whenever possible, each of Indiana's 92 counties.

This report summarizes findings on alcohol, tobacco, marijuana, opioid (prescription-type and illegal), and stimulant use/misuse. In addition, we included data on mental health and suicide, since both substance use and mental distress are highly correlated and frequently co-occur. These data come from a variety of sources,

including national and Indiana-based surveys as well as de-identified administrative records.

As with our prior reports, our primary aim in preparing this annual document is to provide a useful reference tool for policymakers, communities, and professionals involved in substance use prevention and mental health promotion. We realize not everyone has the time or energy to review the contents in detail. For this reason, we again are offering drug fact sheets with summaries on each of the major substances. The full report, as well as earlier versions and supplemental resources, are available on the Center for Health Policy website (<https://fsph.iupui.edu/research-centers/centers/health-policy>). The website also has links to a series of issue briefs related to drug misuse and other behavioral health topics; these briefs are developed each year as part of the SEOW's work. Furthermore, in 2018 we added a Data Portal; i.e., an online tool that allows users to review and interact with data tables, graphs, and maps.

We appreciate your interest and leadership in addressing the problem of substance misuse in Indiana, and, as always, we welcome your feedback on this report and our work.

Marion S. Greene, PhD, MPH

Chair, Indiana State Epidemiological Outcomes Workgroup (SEOW)
Assistant Professor, Health Policy & Management
Center for Health Policy
IU Richard M. Fairbanks School of Public Health at IUPUI
Phone: (317) 278-3247
E-mail: msgreene@iu.edu

1

EXECUTIVE SUMMARY

Substance use continues to be a major public health concern, negatively impacting a variety of health, legal, and social outcomes. Nearly one-fourth of Hoosiers ages 12 and older engaged in binge drinking in the past month and one-tenth used an illicit substance. Furthermore, 7% of Indiana residents met criteria for substance use disorder (SUD) in the past year and 6.5% needed but did not receive treatment for their SUD (Substance Abuse and Mental Health Services Administration [SAMHSA], 2017).

Of particular concern is polysubstance use, i.e., the use of two or more substances over a defined period, simultaneously or at differing times, for recreational purposes. In over 70% of admissions to substance use treatment in Indiana, the use of multiple substances was indicated, with 30% reporting the use of two drugs and nearly 41% reporting the use of three drugs (Indiana Family and Social Services Administration [FSSA], 2018).

During state fiscal year 2018, a total of 10,483 children were removed from their parents by the Department of Child Services in Indiana; almost two-thirds (64%) of these removals were due to parental alcohol and/or drug use (Indiana Department of Child Services, 2018).

Alcohol

Alcohol is the most frequently used substance in Indiana and the United States. Just under half of the population ages 12 and older consumed alcohol within the past month (SAMHSA, 2017). Indiana and U.S. rates of underage drinking among 12- to 17-year-olds were similar (IN: 9.2%; U.S.: 9.6%).

Excessive alcohol consumption contributes to a number of health and economic consequences. Prolonged and compulsive use of alcohol can lead to alcohol use disorder. In 2017, almost one-fourth of Indiana residents ages 12 or older reported binge drinking, which was similar to the national rate (IN: 23.1%; U.S.: 24.4%). About 5% of Hoosiers suffered from alcohol use disorder (AUD) within the past year (U.S.: 5.5%). The highest rate of AUD was found among

18- to 25-year-olds (IN: 10.7%; U.S.: 10.4%) (SAMHSA, 2017).

Alcohol-related collisions decreased from 13,911 in 2003 to 7,733 in 2017. The number of fatal crashes also decreased from 242 to 158 (Indiana State Police, 2017). However, age-adjusted mortality rates for alcohol-attributable deaths have climbed gradually from 2000 through 2017 in both Indiana and the United States. Indiana's age-adjusted rate was 9.6 per 100,000 in 2017, which is the same as the national rate (Centers for Disease Control and Prevention [CDC], 1999-2017).

In addition to health consequences and mortality, alcohol misuse has disproportionately contributed to the United States' economic burden. In 2010, excessive alcohol consumption cost the United States \$249 billion, with Indiana attributing \$4.5 billion (CDC, 2017).

Tobacco / Nicotine

Even though cigarette smoking has declined in recent years, tobacco use is still a public health issue. Cigarette smoking and tobacco-related diseases cost the United States more than \$300 billion per year (CDC, 2018b). In 2017, nearly 22% of Indiana adults reported smoking cigarettes (down from almost 26% in 2011). Indiana's smoking rates have been consistently higher than the national rates (CDC, 2018a).

The decline of cigarette smoking has given rise to other tobacco products. E-cigarettes, hookahs, and other tobacco products gained more popularity and market themselves as safer than cigarettes (Indiana State Department of Health, Tobacco Prevention and Cessation Commission [ISDH/TPCC], 2015). Approximately 23.7% of adults in Indiana reported trying an e-cigarette in 2017 (ISDH/TPCC, 2018). E-cigarettes appeal particularly to younger people and are currently the most commonly used tobacco product among youth. About 24% of Indiana high school students and 22% of Indiana college students reported current use of e-cigarettes (CDC, 1991-2015; King & Jun, 2018).

Tobacco is the leading cause of preventable disease and death in the United States. Tobacco causes 6 million deaths worldwide, about 600,000 of which

are from secondhand smoke exposure (World Health Organization, 2015). The U.S. experiences more than 480,000 deaths from tobacco use, about 41,000 of which are from secondhand smoke exposure (CDC, 2018b). In Indiana, more than 11,100 adults die every year from smoking, and 333,000 live with a tobacco-related disease (U.S. Department of Health and Human Services [USDHHS], 2014).

Opioids

Opioid misuse and addiction have created a national crisis in the United States. According to 2017 data from the National Survey on Drug Use and Health (NSDUH), almost 5% of Indiana residents ages 12 and older misused prescription pain relievers (U.S.: 4.2%) and 0.6% reported using heroin in the past year (U.S.: 0.3%) (SAMHSA, 2017). Rates were generally higher among young adults ages 18 to 25 for both pain relievers (IN: 8.8%; U.S.: 7.1%) and heroin (IN: 1.3%; U.S.: 0.6%) (SAMHSA, 2017).

Opioid treatment programs (OTPs) provide medication-assisted treatment to individuals with opioid use disorder. In Indiana, a total of 13,697 unique patients were treated in OTPs in 2017 (FSSA, 2018). According to the Treatment Episode Data Set (TEDS), in nearly 21% of Indiana treatment admissions, misuse of prescription opioids was reported, and in over 22% of treatment admissions, heroin use was reported (SAMHSA, 2016).

Morbidity and mortality related to opioids increased significantly over the past seven years. Non-fatal emergency department visits due to an opioid overdose rose from 1,856 in 2011 to 8,169 in 2017 (from 45 to 123 visits per 100,000 population) (ISDH, 2018). Overdose deaths involving opioids rose from 347 in 2011 to 1,138 in 2017 (from 5 to 17 deaths per 100,000 population) (ISDH, 2018).

Other Illicit Drugs

Marijuana is the most commonly used illicit drug in the United States (Azofeifa et al., 2016). In 2017, past-month use of marijuana was reported by over 9% of Indiana residents ages 12 and older and past-year use by more than 14%. Prevalence was highest among individuals ages 18 to 25, with nearly 21% of Hoosiers in this age group reporting current marijuana use and 34% reporting past-year use. Indiana and U.S. rates were comparable (SAMHSA, 2017). In nearly half of Indiana treatment admissions, marijuana use was reported (U.S.: 33%) (SAMHSA, 2016).

Stimulants encompass both legal (prescription stimulants such as Ritalin and Adderall) and illicit drugs (such as cocaine and methamphetamine). Almost 2% of Indiana residents ages 12 and older used cocaine in the past year, which was similar to the U.S. rate. Cocaine use was highest among young adults ages 18 to 25, with 6% reporting past year use (SAMHSA, 2017).

Admissions data indicate that methamphetamine was the most widely used stimulant among Hoosiers receiving substance use treatment in 2016. In almost 24% of admissions, methamphetamine use was reported; a percentage significantly higher than the nation's 17%. Cocaine was the second most frequently used stimulant in Indiana's treatment population and was reported in 12% of admissions; a percentage significantly lower than that noted for the rest of the nation (18%). Less than 2% of the treatment population in Indiana and the U.S. reported the use of other stimulants at the time of admission.

REFERENCES, CHAPTER 1

- Azofeifa, A., Mattson, M. E., Schauer, G., McAfee, T., Grant, A., & Lyerla, R. (2016). National Estimates of Marijuana Use and Related Indicators—National Survey on Drug Use and Health, United States, 2002–2014. *MMWR. Surveillance Summaries*, 65, 1-25.
- Centers for Disease Control and Prevention. (1991-2015). Youth Risk Behavior Surveillance System (YRBSS). Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (1999-2017). *CDC WONDER underlying causes of death (compressed mortality)*. Retrieved from <http://wonder.cdc.gov/>
- Centers for Disease Control and Prevention. (2017). *Alcohol and Public Health Data and Maps*. Retrieved from <https://www.cdc.gov/alcohol/data-stats.htm>
- Centers for Disease Control and Prevention. (2018a). *Behavioral Risk Factor Surveillance System (BRFSS) prevalence & trends data*. Retrieved from <http://www.cdc.gov/brfss/brfssprevalence/index.html>
- Centers for Disease Control and Prevention. (2018b). *Burden of Tobacco Use in the U.S.* Retrieved from <https://www.cdc.gov/tobacco/campaign/tips/resources/data/cigarette-smoking-in-united-states.html>
- Indiana Department of Child Services. (2019). SFY2018 *Child Removals due to Parent Substance Abuse*. Source: MaGIK CHINS AFCARS; retrieved 2/1/2019.
- Indiana Family and Social Services Administration. (2018). *Treatment Episode Data System (TEDS)*. Indianapolis, IN: Indiana Family and Social Services Administration.
- Indiana Family and Social Services Administration. (2018). *Opioid Treatment Programs – Patient Count 2017*. Email correspondence with Rhonda Webb from February 15, 2018.
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2015). *Electronic cigarettes*. Retrieved from http://www.in.gov/isdh/tpc/files/Electronic_Cigarettes__12_9_2015.pdf
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2018). Indiana Youth Tobacco Survey and Indiana Adult Tobacco Survey. Data received from Katelin Rupp, Director of Program Evaluation at the Indiana State Department of Health, Tobacco Prevention & Cessation Commission.
- Indiana State Department of Health. (2018). *Stats Explorer*. Retrieved from https://gis.in.gov/apps/isdh/meta/stats_layers.htm
- Indiana State Police. (2017). *Automated Reporting Information Exchange System (ARIES), Vehicle Crash Records System*. Data received from the Center for Criminal Justice Research, Public Policy Institute, School of Public and Environmental Affairs, Indiana University–Purdue University Indianapolis.
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey, 2018*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey>
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2016). *Treatment Episode Data Set -- Admissions (TEDS-A)*. Retrieved from <http://www.dasis.samhsa.gov/dasis2/teds.htm>
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>
- U.S. Department of Health and Human Services. (2014). *The Health Consequences of Smoking -- 50 years of progress: A Report of the Surgeon General*. Atlanta, GA.: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health.

2

ALCOHOL USE IN INDIANA: CONSUMPTION PATTERNS AND CONSEQUENCES

INTRODUCTION

Alcohol is the most frequently used substance in both Indiana and the United States. In 2016, the National Institute on Alcohol Abuse and Alcoholism (NIAAA) estimated that Hoosiers 14 years and older consumed 11,756 gallons of ethanol (the intoxicating agent in alcoholic beverages). By volume, this equates to 118,132 gallons of beer, 12,321 gallons of wine, or 11,802 gallons of spirits. This level of use represents an annual per capita consumption rate of 2.2 gallons of ethanol for Hoosiers age 14 and older (NIAAA, 2018). In 2017, there were 12,999 alcohol beverage permits on file in Indiana, representing a rate of 1.8 licenses per 1,000 Hoosiers; thus, Indiana residents have many points of access to alcohol (Alcohol and Tobacco Commission, 2017).

Alcohol's legal status, its wide availability, and its social acceptability are all contributors to patterns of excessive or risky use, such as heavy drinking or binge drinking. Excessive consumption of alcohol is responsible for significant morbidity and mortality due to alcohol-related health problems (e.g., cirrhosis and other serious liver diseases), alcohol use disorders, homicides, suicides, violent crimes, and vehicle crashes. Additionally, other health-compromising behaviors such as cigarette smoking, illicit drug use, and risky sexual behaviors have also been linked to drinking (CDC, 2016).

Alcohol use can also contribute to adverse social outcomes such as job loss and involvement with the criminal justice and social service system. In 2010, the most recent year for which estimates are available, Indiana spent \$4.5 billion to deal with the negative consequences of excessive alcohol use, with much of these expenses tied to outcomes associated with binge drinking (Sacks, Gonzales, Bouchery, Tomedi, & Brewer, 2015).

PREVALENCE OF ALCOHOL CONSUMPTION IN THE GENERAL POPULATION


National Survey on Drug Use and Health

Based on 2016–2017 averages from the Substance Abuse and Mental Health Services Administration (SAMHSA)'s National Survey on Drug Use and Health (NSDUH), an estimated 49.9% (95% Confidence Interval [CI]: 47.1–52.7) of Indiana residents 12 years of age or older had used alcohol in the past month; Indiana's prevalence rate for current alcohol use¹ was similar to the U.S. rate of 51.2% (95% CI: 50.8–51.7) (see Figure 2.1). Young adults between the ages of 18 and 25 reported the highest level of use. Rates were similar between Indiana residents in that age group (59.8%; 95% CI: 55.8–63.7) and their national counterparts (56.7%, 95% CI: 55.9–57.6). Furthermore, 9.2% (95% CI: 7.6–11.2) of young people ages 12 to 17 consumed alcohol in the past 30 days in Indiana (see Figure 2.2); the rate was similar on the national level (9.6%; 95% CI: 9.1–10.0).

NSDUH also provides underage drinking estimates for 12- to 20-year-olds. In 2017, Indiana's rate for current alcohol use in underage Hoosiers (20.2%; 95% CI: 17.8–22.7) was similar to that of the U.S. (19.5%; 95% CI: 18.8–20.2) (SAMHSA, 2017).


¹ Current alcohol use is defined as having used alcohol in the past 30 days or past month.

Figure 2.1 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Current Alcohol Use (National Survey on Drug Use and Health, 2008–2017)



Source: SAMHSA, 2017

Figure 2.2 Percentage of Indiana Population Reporting Current Alcohol Use by Age Group (National Survey on Drug Use and Health, 2008–2017)




Source: SAMHSA, 2017

In 2015, SAMHSA redesigned the questions on the NSDUH pertaining to binge drinking. The definition of binge drinking for women was lowered from five or more drinks on one occasion to four or more drinks (for men, it remained at five or more drinks). 2016 is the first year for which both national- and state-level estimates are

available. These new estimates of binge drinking cannot be compared with estimates from previous years (Center for Behavioral Health Statistics and Quality, 2016). Based on the new definition for binge drinking, the NSDUH estimated that in 2017, 23.1% of Indiana’s population 12 years of age or older reported binge drinking (95%

Figure 2.3 Current Binge Drinking in Indiana and the U.S. by Age Group (National Survey on Drug Use and Health, 2017)



Source: SAMHSA, 2017

Table 2.1 Percentage of Indiana Adults Having Used Alcohol in the Past 30 Days, by Gender, Race/Ethnicity, and Age Group (Behavioral Risk Factor Surveillance System, 2017)

		Indiana % (95% CI)
Gender	Male	58.7% (57.0–60.4)
	Female	45.0% (43.4–46.6)
Race/Ethnicity	White	51.9% (50.7–53.2)
	Black	50.3% (45.7–54.8)
	Hispanic	50.4% (45.2–55.7)
Age Group	18-24	52.9% (48.4–57.3)
	25-34	63.8% (60.5–67.0)
	35-44	58.4% (55.3–61.5)
	45-54	53.8% (51.2–56.4)
	55-64	48.9% (46.7–51.1)
	65+	36.5% (34.9–38.1)
Total		51.6% (50.5–52.8)


Source: CDC, 2018

CI: 20.8–25.5); this represents a rate similar to the national average of 24.4% (95% CI: 24.0–24.8). Binge drinking was more prevalent among 18- to 25-year-olds than among any other age group (U.S.: 37.6%; 95% CI: 36.9–38.4; IN: 38.2%; 95% CI: 34.3–42.4). Binge drinking rates in 2017 in underaged individuals were similar in Indiana (12.5%; 95% CI: 10.7–14.5) and the U.S. (12.0%; 95% CI: 11.5–12.5) (SAMHSA, 2017) (see Figure 2.3).

Behavioral Risk Factor Surveillance System

Based on findings from the Centers for Disease Control and Prevention (CDC)'s Behavioral Risk Factor Surveillance System (BRFSS), adult prevalence rates for current alcohol use in 2017 were 51.6% (95% CI: 50.5–52.8) for Indiana and 54.7% for the nation. In Indiana, rates tended to be higher among males and among younger age groups (see Table 2.1) (CDC, 2018).

Figure 2.4 Percentage of Indiana and U.S. Adults Reporting Binge Drinking in the Past 30 Days (Behavioral Risk Factor Surveillance System, 2011–2017)



Source: CDC, 2018

Table 2.2 Percentage of Indiana Residents Who Engaged in Binge Drinking in the Past 30 Days, by Gender, Race/Ethnicity, and Age Group (Behavioral Risk Factor Surveillance System, 2017)

		Indiana % (95% CI)
Gender	Male	22.3% (20.8–23.8)
	Female	11.2% (10.1–12.3)
Race/Ethnicity	White	16.4% (15.4–17.4)
	Black	14.9% (11.7–18.2)
	Hispanic	21.6% (16.9–26.2)
Age Group	18-24	24.4% (20.8–28.1)
	25-34	27.1% (24.1–30.1)
	35-44	19.7% (17.2–22.2)
	45-54	17.6% (15.6–19.6)
	55-64	10.5% (9.1–11.8)
	65+	5.0% (4.3–5.8)
Total		16.6% (15.7–17.5)

Source: CDC, 2017

The BRFSS defines binge drinking as “males having five or more drinks on one occasion and females having four or more drinks on one occasion.” The overall prevalence rate for adult binge drinking in Indiana (16.6%, 95% CI: 15.7–17.5) was similar to the U.S. median rate (17.4%) in 2017.

Statewide, binge alcohol use was significantly higher in males and more prevalent in younger individuals (see Table 2.2). Trends in binge drinking are shown in Figure 2.4 (CDC, 2018).

Youth Risk Behavior Surveillance System

According to the CDC’s Youth Risk Behavior Surveillance System (YRBSS), in 2015, 30.5% (95% CI: 26.3–35.2) of Indiana high school students had consumed at least one alcoholic drink in the past 30 days. No significant differences in alcohol consumption were observed by gender or race/ethnicity; however, rates varied by grade level, with 9th grade students reporting the lowest rate. Indiana’s past-month alcohol prevalence among high school students was similar to the nation’s rate (32.8%: 95% CI: 30.4–35.2). Furthermore, 17.4% (95% CI: 14.0–21.5) of Indiana high school students reported having had


five or more alcoholic drinks within a couple of hours at least once in the past month; the U.S. rate was similar at 17.7% (95% CI: 15.8–19.8). Indiana’s binge alcohol consumption among high school students decreased significantly from 28.9% in 2003 to 17.4% in 2015 (CDC, 1991–2015).

Indiana Youth Survey

The Indiana Youth Survey (INYS) indicates that in 2018, 29.5% of Indiana 12th grade students reported using


alcohol at least once during the past 30 days (Gassman et al., 2018). Overall, alcohol consumption patterns seemed to progress with age; i.e., 8th grade students showed lower prevalence rates than 10th and 12th grade students. For more detailed data on monthly alcohol use among Indiana and U.S. 8th, 10th, and 12th grade students, see Figure 2.5; for trend information (from 2009 through 2018) on monthly alcohol use among high school seniors, see Figure 2.6. For monthly and binge use by Indiana region and grade for 2018, see Appendix 2A.

Figure 2.5 Percentage of Indiana and U.S. 8th, 10th, and 12th Grade Students Reporting Monthly Alcohol Use (Indiana Youth Survey and Monitoring the Future Survey, 2018)



Source: Gassman et al., 2018; Inter-university Consortium for Political and Social Research, University of Michigan, 2018

Figure 2.6 Percentage of Indiana and U.S. High School Seniors (12th Grade) Reporting Monthly Alcohol Use (Indiana Youth Survey and Monitoring the Future Survey, 2009–2018)



Source: Gassman et al., 2018; Inter-university Consortium for Political and Social Research, University of Michigan, 2018

Indiana College Substance Use Survey

The Indiana Prevention Resource Center (IPRC) developed the Indiana College Substance Use Survey (ICSUS) to measure alcohol and other drug usage, attitudes, and perceptions among college students at two- and four-year institutions. According to 2018 results, 63.3% of respondents reported past-month alcohol use; past-month consumption rates were significantly lower for underage students (53.1%) than for those ages 21 and older (80.0%). Similarly, past-month binge drinking prevalence (overall 36.7%) was significantly lower for underage students (31.0%) than for those ages 21 and older (46.2%) (King & Jun, 2018).²

USE OF ALCOHOL IN THE TREATMENT POPULATION National Survey on Drug Use and Health

Based on 2016–2017 NSDUH averages, the estimated prevalence for alcohol use disorder³ in the past year among those ages 12 and older was 5.2% (95% CI: 4.3–6.3) in Indiana, which was similar to the national estimate (5.5%; 95% CI: 5.3–5.6) (see Figure 2.7). Of all age groups, adults ages 18 to 25 reported the highest prevalence rates both in Indiana and nationally across all years reviewed. Additionally, an estimated 4.9% (95% CI: 4.1–5.9) of those ages 12 and older were in need of but did not receive treatment for alcohol use in Indiana (U.S.: 5.2%; 95% CI: 5.1–5.4) (SAMHSA, 2017).

²Twenty-four (24) Indiana colleges participated in the 2018 survey; results are based on nonrandom sampling and are not representative of all college students in Indiana.


³The NSDUH defines alcohol use disorder as meeting the criteria for “dependence” or “abuse” based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Treatment Episode Data Set

According to the Treatment Episode Data Set (TEDS), alcohol plays a major role in admissions to substance abuse treatment. In nearly half (47.8%) of Indiana


treatment episodes in 2016, alcohol use was reported (U.S.: 46.9%), and in 29.6%, alcohol dependence⁴ was indicated (U.S.: 32.6%) (see Figure 2.8) (SAMHSA, 2015).

Figure 2.7 Percentage of Indiana and U.S. Population Ages 12 and Older with Alcohol Use Disorder (National Survey on Drug Use and Health, 2008–2017)



Source: SAMHSA, 2017

Figure 2.8 Percentage of Treatment Episodes in Indiana and the United States with Alcohol Use and Alcohol Dependence Reported at Treatment Admission (Treatment Episode Data Set, 2007–2016)



Source: SAMHSA, 2016

⁴We defined alcohol dependence as “individuals in substance abuse treatment listing alcohol as their primary substance at admission.”

Factors significantly associated with alcohol use in Indiana’s treatment population included gender, race/ethnicity, and age:

Gender—A higher percentage of males (53.8%) in substance abuse treatment reported alcohol use, compared to 41.4% of females.

Race/ethnicity—Nearly half (45.5%) of whites in treatment reported using alcohol at the time of admission; this percentage was higher for blacks (59.5%) and other races (52.5%). With regard to ethnicity, a significantly higher percentage of Hispanics (61.6%) reported alcohol use than non-Hispanics (47.2%).

Age—The percentage of Hoosiers reporting alcohol use at treatment admission increased with age and was highest among those ages 55 and older (74.5%).

Table 2.3 depicts the percentage of Indiana residents, categorized by gender, race, ethnicity, and age group, reporting alcohol use at treatment admission. See Appendix 2B, for county-level treatment data.

Table 2.3 Percentage of Treatment Episodes in Indiana Reporting Alcohol Use at Treatment Admission, by Gender, Race, Ethnicity, and Age Group (Treatment Episode Data Set, 2016)

		Alcohol Dependence
Gender	Male	53.8%
	Female	38.9%
Race	White	45.5%
	Black	59.5%
	Other	52.5%
Ethnicity	Hispanic	61.6%
	Non-Hispanic	47.2%
Age Group	Under 18	42.1%
	18-24	39.9%
	25-34	39.3%
	35-44	51.0%
	45-54	68.2%
	55+	74.5%
Total		47.8%

Source: SAMHSA, 2016

CONSEQUENCES OF ALCOHOL USE

Hospitalizations

Hospital discharge records show that in 2017, a total of 7,116 hospitalized patients were treated in Indiana for an alcohol-attributable primary diagnosis, representing one percent (0.9%) of all hospital discharges in the state (Indiana State Department of Health [ISDH], 2017).⁵

Fetal Alcohol Spectrum Disorders

Alcohol consumption during pregnancy is another major health concern since fetal alcohol spectrum disorders (FASD) are a direct result of prenatal exposure to alcohol. FASD is not a clinical diagnosis, but an umbrella term used to describe a range of disorders such as fetal alcohol syndrome, alcohol-related neurodevelopmental disorder, and alcohol-related birth defects. Possible

physical effects include brain damage; facial anomalies; growth deficiencies; defects of heart, kidney, and liver; vision and hearing problems; skeletal defects; and dental abnormalities. In the United States, the prevalence of fetal alcohol spectrum disorders is 10.0 per 1,000 live births (SAMHSA, Fetal Alcohol Spectrum Disorders Center for Excellence, 2007).

The Indiana Birth Defects and Problems Registry collects information on birth defects and birth problems for all children in Indiana from birth to 3 years old (5 years old for autism and fetal alcohol syndrome). State law requires doctors, hospitals, and other healthcare providers to submit a report to the registry at ISDH when a child is born with a birth defect. From 2015 through 2017, 61 children were born with fetal alcohol syndrome,⁶ the most severe form of FASD, in Indiana (ISDH, 2015–2017).

⁵For our analysis, we only included primary diagnoses that were 100% attributable to alcohol, as listed in CDC’s Alcohol-Related Disease Impact (ARDI) database. These included ICD-10 codes E24.4 (Alcohol-induced pseudo-Cushing’s syndrome), F10 (Mental and behavioral disorders due to use of alcohol), G31.2 (Degeneration of nervous system due to alcohol), G62.1 (Alcoholic polyneuropathy), G72.1 (Alcoholic myopathy), I42.6 (Alcoholic cardiomyopathy), K29.2 (Alcoholic gastritis), K70 (Alcoholic liver disease), K86.0 (Alcohol-induced chronic pancreatitis), R78.0 (Finding of alcohol in blood), X45 (Accidental poisoning by and exposure to alcohol), X65 (Intentional self-poisoning by and exposure to alcohol), Y15 (Poisoning by and exposure to alcohol, undetermined cause) (CDC, 2006-2010).


⁶The ICD-9 code for fetal alcohol syndrome is 760.71.

Alcohol-Related Mortality

From 2000 through 2016, a total of 8,034 Hoosiers died from alcohol-induced causes, and mortality rates attributable to alcohol have climbed gradually in both Indiana and the United States (CDC, 1999–2017).⁷ In

2017, Indiana’s age-adjusted alcohol-attributable death rate was 9.6 per 100,000 (95% CI: 8.9–10.3); the same as the U.S. rate (95% CI: 9.5–9.7) (see Figure 2.9) (CDC, 1999–2017).

Figure 2.9 Age-Adjusted Alcohol-attributable Mortality Rates per 100,000 Population in Indiana and the United States (CDC WONDER, 2008–2017)



Source: CDC, 1999–2017

⁷Alcohol-induced causes of death include the following ICD-10 codes: E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K86.0, R78.0, X45, X65, Y15.

Appendix 2C, lists conditions that can be attributed to alcohol, along with their alcohol-attributable percentages. The list was developed through CDC's Alcohol-Related Disease Impact (ARDI) database (CDC, 2006–2010).

Alcohol-Related Motor Vehicle Accidents

Data from the Automated Reporting Information Exchange System (ARIES), part of the Indiana State Police's Vehicle Crash Records System, showed a decrease in alcohol-related collisions from 13,911 in 2003 to 7,733 in 2017. This represents a 44% drop. The number of fatal crashes with alcohol involvement also decreased, from 242 to 158, representing a 35% drop. (For a detailed listing of alcohol-related collisions and fatalities in Indiana by county for 2017, see Appendix 2D). The overall rate for alcohol-related collisions in Indiana in 2017 was 1.2 per 1,000 population (Indiana State Police, 2017).

Child Removals due to Parental Substance Abuse

During SFY 2018, a total of 10,483 children were removed from their homes. In 847 cases (8.1%), parental alcohol abuse was indicated as the reason for removal (Indiana Department of Child Services, 2019).⁸ [See Appendix 2E for county-level information.]

Alcohol, Tobacco, and/or Drug-Related School Suspensions or Expulsions

In Indiana, students can be suspended or expelled from school for using alcohol, tobacco, and/or drugs on school property. Data from the Indiana Department of Education (IDOE) indicate that during the academic year 2018, a total of 1,006 suspensions/expulsions were recorded in Indiana schools related to alcohol (IDOE, 2019). [See Appendix 2F for county-level information.]

⁸Counts and percentages may underrepresent removals that involve parental alcohol and/or drug abuse as data relies on parent alcohol and/or drug abuse being selected as a removal reason. There may be instances where alcohol and/or drug abuse is present but not selected as the removal reason.

APPENDIX 2A

Percentage of Indiana Students Reporting Monthly and Binge Alcohol Use, by Region and Grade (Indiana Youth Survey, 2018)

		Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
6th Grade	Monthly	4.0	4.6	3.7	2.8	3.6	3.9	4.1	3.2	4.6
	Binge	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
7th Grade	Monthly	7.6	7.6	6.6	6.1	5.8*	9.3*	7.7	6.3*	9.1*
	Binge	2.5	2.4	1.9	2.1	1.9	3.5*	2.0	2.1	3.5*
8th Grade	Monthly	13.0	15.1*	12.3	15.7	10.0*	13.1	12.4	12.0	13.9
	Binge	4.7	4.8	4.4	6.4	3.5*	5.2	4.7	4.5	5.1
9th Grade	Monthly	16.3	19.1*	17.0	16.4	16.2	12.0*	16.0	16.9	15.5
	Binge	6.2	7.0	6.6	5.9	4.6*	4.7*	5.6	7.6*	6.4
10th Grade	Monthly	21.0	24.2*	17.2*	18.2*	20.4	18.4*	20.7	23.9*	21.8
	Binge	8.5	9.7*	6.7*	7.2	8.3	8.1	8.0	10.9*	8.3
11th Grade	Monthly	24.1	24.7	21.5*	21.3	24.0	22.0	23.5	28.3*	24.5
	Binge	10.8	10.5	8.6*	9.1	9.9	9.4	9.2	15.5*	11.6
12th Grade	Monthly	29.5	30.2	24.8*	22.6*	29.0	29.6	25.5*	33.0*	33.2*
	Binge	13.8	13.8	10.9*	8.4*	14.6	15.1	10.2*	16.8*	15.9*

Notes: * Indicates a local rate that is significantly different from the overall state rate (P < 0.05).
 Beginning in 2015, the Indiana Youth Survey stopped asking 6th grade students about binge drinking.
 Source: Gassman et al., 2018

APPENDIX 2B

Number of Treatment Episodes with Alcohol Use and Dependence Reported at Treatment Admission in Indiana, by County (Substance Abuse Population by County/Treatment Episode Data Set, 2018)

County	Treatment Episodes	Alcohol Use		Alcohol Dependence	
	Total	Number	%	Number	%
Adams	228	126	55.3%	78	34.2%
Allen	2,050	1213	59.2%	709	34.6%
Bartholomew	727	218	30.0%	119	16.4%
Benton	61	38	62.3%	26	42.6%
Blackford	144	38	26.4%	13	9.0%
Boone	167	70	41.9%	38	22.8%
Brown	72	27	37.5%	15	20.8%
Carroll	80	40	50.0%	25	31.3%
Cass	387	179	46.3%	110	28.4%
Clark	103	27	26.2%	23	22.3%
Clay	106	38	35.8%	29	27.4%
Clinton	147	67	45.6%	38	25.9%
Crawford	36	16	44.4%	7	19.4%
Daviess	202	93	46.0%	63	31.2%
Dearborn	497	240	48.3%	142	28.6%
Decatur	185	77	41.6%	50	27.0%
DeKalb	276	160	58.0%	93	33.7%
Delaware	843	264	31.3%	190	22.5%
Dubois	131	70	53.4%	37	28.2%
Elkhart	765	398	52.0%	246	32.2%
Fayette	322	73	22.7%	34	10.6%
Floyd	49	9	18.4%	7	14.3%
Fountain	59	24	40.7%	10	16.9%
Franklin	145	52	35.9%	27	18.6%
Fulton	188	103	54.8%	53	28.2%
Gibson	252	146	57.9%	82	32.5%
Grant	509	200	39.3%	114	22.4%
Greene	197	80	40.6%	43	21.8%
Hamilton	874	556	63.6%	339	38.8%
Hancock	398	174	43.7%	132	33.2%
Harrison	19	6	31.6%	<5	N/A
Hendricks	424	204	48.1%	120	28.3%
Henry	381	119	31.2%	77	20.2%
Howard	492	203	41.3%	77	15.7%
Huntington	136	51	37.5%	29	21.3%
Jackson	330	151	45.8%	71	21.5%
Jasper	93	33	35.5%	21	22.6%
Jay	122	31	25.4%	18	14.8%
Jefferson	307	96	31.3%	52	16.9%
Jennings	260	90	34.6%	57	21.9%
Johnson	424	148	34.9%	102	24.1%
Knox	352	177	50.3%	112	31.8%
Kosciusko	362	213	58.8%	103	28.5%
LaGrange	115	58	50.4%	32	27.8%
Lake	1,903	1,098	57.7%	813	42.7%
LaPorte	172	72	41.9%	48	27.9%
Lawrence	535	201	37.6%	91	17.0%
Madison	1,205	489	40.6%	279	23.2%
Marion	4,134	1,841	44.5%	1267	30.6%
Marshall	137	59	43.1%	41	29.9%
Martin	60	30	50.0%	24	40.0%
Miami	249	118	47.4%	70	28.1%
Monroe	1,428	689	48.2%	423	29.6%
Montgomery	304	110	36.2%	59	19.4%
Morgan	550	205	37.3%	124	22.5%
Newton	27	13	48.1%	5	18.5%
Noble	346	173	50.0%	115	33.2%
Ohio	33	21	63.6%	12	36.4%
Orange	135	57	42.2%	37	27.4%
Owen	187	95	50.8%	59	31.6%
Parke	65	22	33.8%	10	15.4%
Perry	94	50	53.2%	29	30.9%
Pike	46	20	43.5%	18	39.1%
Porter	202	81	40.1%	50	24.8%
Posey	156	90	57.7%	48	30.8%
Pulaski	82	46	56.1%	32	39.0%
Putnam	253	93	36.8%	37	14.6%
Randolph	167	44	26.3%	28	16.8%
Ripley	180	94	52.2%	54	30.0%
Rush	139	44	31.7%	24	17.3%
Saint Joseph	1,351	658	48.7%	430	31.8%
Scott	212	56	26.4%	33	15.6%
Shelby	140	61	43.6%	46	32.9%
Spencer	132	61	46.2%	33	25.0%
Starke	77	19	24.7%	14	18.2%
Steuben	183	116	63.4%	63	34.4%
Sullivan	115	61	53.0%	34	29.6%
Switzerland	113	50	44.2%	29	25.7%
Tippecanoe	364	160	44.0%	82	22.5%
Tipton	46	18	39.1%	9	19.6%
Union	54	21	38.9%	17	31.5%
Vanderburgh	1,218	580	47.6%	317	26.0%
Vermillion	132	63	47.7%	37	28.0%
Vigo	771	311	40.3%	179	23.2%
Wabash	299	129	43.1%	64	21.4%
Warren	26	11	42.3%	7	26.9%
Warrick	310	166	53.5%	77	24.8%
Washington	44	14	31.8%	10	22.7%
Wayne	558	209	37.5%	125	22.4%
Wells	156	70	44.9%	30	19.2%
White	125	63	50.4%	28	22.4%
Whitley	126	81	64.3%	45	35.7%
Indiana	18,129	15,229	45.7%	9,272	27.8%

Notes: We defined alcohol dependence as “individuals in substance abuse treatment listing alcohol as their primary substance at admission.”

We calculated the percentages by dividing the number of reported alcohol use/dependence by the number of treatment episodes.

Information on treatment episodes <5 was suppressed due to confidentiality constraints.

Source: Indiana Family and Social Services Administration, 2018

APPENDIX 2C

Conditions that are Directly Attributable to Alcohol in Indiana (Alcohol-Related Disease Impact, Based on Averages from 2006–2010)

Condition	Percentage Directly Attributable to Alcohol	Condition	Percentage Directly Attributable to Alcohol
Alcohol abuse/dependence	100%	Chronic pancreatitis	84%
Alcohol cardiomyopathy	100%	Gastroesophageal hemorrhage	47%
Alcohol polyneuropathy	100%	Homicide	47%
Alcohol-induced chronic pancreatitis	100%	Fire Injuries	42%
Alcoholic gastritis	100%	Hypothermia	42%
Alcoholic liver disease	100%	Esophageal varices	40%
Alcoholic myopathy	100%	Liver cirrhosis, unspecified	40%
Alcoholic psychosis	100%	Portal hypertension	40%
Degeneration of nervous system due to alcohol	100%	Drowning	34%
Fetal alcohol syndrome/Fetus and newborn affected by maternal alcohol use	100%	Fall injuries	32%
Alcohol poisoning	100%	Poisoning (not alcohol)	29%
Excessive blood alcohol level	100%	Acute pancreatitis	24%
Suicide by and exposure to alcohol	100%	Suicide	23%

Source: Centers for Disease Control and Prevention, 2006–2010

APPENDIX 2D

Number and Rate (per 1,000) of All and Fatal Alcohol-Related Collisions in Indiana, by County (Automated Reporting Information Exchange System, 2017)

County	All Collisions			Fatal Collisions		
	Total Collisions	Alcohol-related Collisions	Alcohol-related Collision Rate	Total Fatal Collision	Alcohol-related Fatal Collisions	Alcohol-related Fatal Collision Rate
Adams	741	27	0.76	8	1	0.03
Allen	14,123	541	1.45	40	15	0.04
Bartholomew	2,054	61	0.74	15	2	0.02
Benton	159	11	1.28	1	0	0.00
Blackford	296	12	1.00	1	1	0.08
Boone	2,058	66	1.00	7	0	0.00
Brown	597	34	2.26	7	1	0.07
Carroll	527	27	1.35	2	0	0.00
Cass	1,293	48	1.26	7	0	0.00
Clark	4,496	103	0.88	14	0	0.00
Clay	696	27	1.03	4	0	0.00
Clinton	1,164	44	1.36	11	2	0.06
Crawford	373	15	1.42	2	1	0.09
Daviess	320	29	0.88	6	3	0.09
Dearborn	1,824	66	1.33	7	1	0.02
Decatur	874	38	1.42	1	0	0.00
DeKalb	1,385	51	1.19	4	1	0.02
Delaware	4,102	152	1.32	8	1	0.01
Dubois	1,553	70	1.64	5	1	0.02
Elkhart	7,509	197	0.96	24	4	0.02
Fayette	506	18	0.78	4	1	0.04
Floyd	2,839	86	1.12	10	4	0.05
Fountain	443	27	1.64	1	0	0.00
Franklin	513	29	1.28	4	1	0.04

(Continued on next page)

APPENDIX 2D (Continued from previous page)

County	All Collisions			Fatal Collisions		
	Total Collisions	Alcohol-related Collisions	Alcohol-related Collision Rate	Total Fatal Collision	Alcohol-related Fatal Collisions	Alcohol-related Fatal Collision Rate
Fulton	631	20	1.00	3	1	0.05
Gibson	1,084	37	1.10	16	2	0.06
Grant	2,331	86	1.29	8	3	0.05
Greene	925	26	0.81	11	2	0.06
Hamilton	8,812	230	0.71	10	3	0.01
Hancock	1,850	53	0.71	15	5	0.07
Harrison	1,316	41	1.03	9	1	0.03
Hendricks	4,573	143	0.87	8	0	0.00
Henry	1,037	47	0.97	4	1	0.02
Howard	2,654	91	1.10	11	6	0.07
Huntington	1,190	35	0.96	4	1	0.03
Jackson	1,715	46	1.05	10	1	0.02
Jasper	1,221	52	1.55	7	0	0.00
Jay	577	19	0.91	3	0	0.00
Jefferson	1,049	34	1.06	8	2	0.06
Jennings	774	31	1.12	10	3	0.11
Johnson	3,650	138	0.90	12	0	0.00
Knox	930	49	1.31	7	4	0.11
Kosciusko	2,750	95	1.20	14	0	0.00
LaGrange	1,027	46	1.17	9	0	0.00
Lake	17,419	698	1.44	49	16	0.03
LaPorte	3,814	194	1.76	21	5	0.05
Lawrence	1,553	51	1.12	12	0	0.00
Madison	4,189	156	1.20	13	4	0.03
Marion	36,963	1,181	1.24	98	16	0.02
Marshall	1,588	57	1.23	6	0	0.00
Martin	119	4	0.39	2	0	0.00
Miami	1,076	45	1.26	7	0	0.00
Monroe	4,297	161	1.10	8	0	0.00
Montgomery	1,123	35	0.91	4	0	0.00
Morgan	1,943	86	1.23	4	0	0.00
Newton	383	32	2.26	6	3	0.21
Noble	1,393	56	1.18	7	0	0.00
Ohio	183	8	1.37	1	1	0.17
Orange	610	22	1.13	3	0	0.00
Owen	603	26	1.25	6	0	0.00
Parke	461	14	0.83	2	1	0.06
Perry	500	26	1.36	1	0	0.00
Pike	137	17	1.37	1	0	0.00
Porter	5,143	224	1.33	25	1	0.01
Posey	585	28	1.09	3	0	0.00
Pulaski	446	8	0.64	4	0	0.00
Putnam	1,035	38	1.01	8	2	0.05
Randolph	520	20	0.80	7	0	0.00
Ripley	822	31	1.09	5	2	0.07
Rush	347	12	0.72	0	0	0.00
Saint Joseph	9,041	295	1.09	24	4	0.01
Scott	628	23	0.96	5	1	0.04
Shelby	1,307	57	1.28	7	0	0.00
Spencer	565	27	1.32	6	1	0.05

(Continued on next page)

APPENDIX 2D (Continued from previous page)

County	All Collisions			Fatal Collisions		
	Total Collisions	Alcohol-related Collisions	Alcohol-related Collision Rate	Total Fatal Collision	Alcohol-related Fatal Collisions	Alcohol-related Fatal Collision Rate
Starke	622	19	0.83	3	1	0.04
Steuben	1,526	55	1.59	5	1	0.03
Sullivan	451	21	1.01	2	2	0.10
Switzerland	124	5	0.47	2	1	0.09
Tippecanoe	7,275	213	1.12	19	6	0.03
Tipton	383	21	1.39	4	2	0.13
Union	86	8	1.11	2	0	0.00
Vanderburgh	6,834	205	1.13	19	3	0.02
Vermillion	336	16	1.03	5	1	0.06
Vigo	3,799	134	1.25	13	3	0.03
Wabash	871	41	1.30	1	0	0.00
Warren	233	13	1.59	3	1	0.12
Warrick	1,575	55	0.88	5	1	0.02
Washington	671	37	1.33	11	2	0.07
Wayne	2,314	70	1.06	7	1	0.02
Wells	752	22	0.79	4	0	0.00
White	900	31	1.28	5	0	0.00
Whitley	1,019	37	1.10	7	1	0.03
Indiana	219,105	7,733	1.16	834	158	0.02

Note: Rates based on numbers lower than 20 are unreliable.
Source: Indiana State Police, 2017

APPENDIX 2E

Child Removals, Total and Due to Parental Alcohol Abuse, SFY 2018

County	Removals Total	Parent Alcohol Abuse Indicated as Removal Reason		County	Removals Total	Parent Alcohol Abuse Indicated as Removal Reason	
		Count	Percentage			Count	Percentage
Adams	66	11	16.7	Madison	192	18	9.4
Allen	450	38	8.4	Marion	1775	103	5.8
Bartholomew	104	4	3.8	Marshall	80	10	12.5
Benton	9	1	11.1	Martin	16	0	0
Blackford	32	0	0	Miami	57	4	7.0
Boone	54	5	9.3	Monroe	158	24	15.2
Brown	40	1	2.5	Montgomery	100	10	10.0
Carroll	31	1	3.2	Morgan	133	15	11.3
Cass	41	0	0	Newton	17	3	17.6
Clark	126	10	7.9	Noble	75	7	9.3
Clay	78	13	16.7	Ohio	7	0	0
Clinton	57	3	5.3	Orange	35	2	5.7
Crawford	23	2	8.7	Owen	52	8	15.4
Daviess	60	10	16.7	Parke	35	4	11.4
Dearborn	82	7	8.5	Perry	70	14	20.0
Decatur	77	14	18.2	Pike	46	1	2.2
Dekalb	37	2	5.4	Porter	129	14	10.9
Delaware	230	18	7.8	Posey	85	2	2.4
Dubois	81	3	3.7	Pulaski	27	2	7.4
Elkhart	169	20	11.8	Putnam	72	2	2.8
Fayette	58	4	6.9	Randolph	56	7	12.5
Floyd	161	9	5.6	Ripley	98	6	6.1
Fountain	45	5	11.1	Rush	14	0	0
Franklin	29	1	3.4	St. Joseph	353	32	9.1
Fulton	71	4	5.6	Scott	126	12	9.5
Gibson	140	12	8.6	Shelby	59	2	3.4
Grant	92	7	7.6	Spencer	64	3	4.7
Greene	73	7	9.6	Starke	68	0	0
Hamilton	86	9	10.5	Steuben	52	6	11.5
Hancock	78	8	10.3	Sullivan	71	5	7.0
Harrison	35	2	5.7	Switzerland	23	0	0
Hendricks	101	17	16.8	Tippecanoe	225	16	7.1
Henry	110	9	8.2	Tipton	39	0	0
Howard	96	7	7.3	Union	7	0	0
Huntington	38	4	10.5	Vanderburgh	595	46	7.7
Jackson	73	13	17.8	Vermillion	57	2	3.5
Jasper	69	2	2.9	Vigo	263	17	6.5
Jay	35	0	0	Wabash	65	5	7.7
Jefferson	79	5	6.3	Warren	18	1	5.6
Jennings	83	1	1.2	Warrick	107	3	2.8
Johnson	114	7	6.1	Washington	27	0	0
Knox	115	18	15.7	Wayne	117	9	7.7
Kosciusko	104	6	5.8	Wells	54	6	11.1
LaGrange	38	3	7.9	White	53	5	9.4
Lake	684	69	10.1	Whitley	42	14	33.3
Laporte	102	8	7.8	Indiana	10,483	847	8.1%
Lawrence	105	6	5.7				

Note: Counts and percentages may underrepresent removals that involve parental alcohol and/or drug abuse as data relies on parent alcohol and/or drug abuse being selected as a removal reason. There may be instances where alcohol and/or drug abuse is present but not selected as the removal reason.

Source: Indiana Department of Child Services, 2018

APPENDIX 2F

School Suspensions or Expulsions Related to Alcohol, Tobacco, and/or Drug Use (2018)

County	Total	Count	Percentage	County	Total	Count	Percentage
Adams	4,347	<5	<5	Madison	20,089	16	15
Allen	57,046	68	68	Marion	179,578	119	112
Bartholomew	13,126	16	16	Marshall	7,759	19	16
Benton	1,928	<5	<5	Martin	1,443	<5	<5
Blackford	1,764	<5	<5	Miami	7,480	10	10
Boone	12,342	6	6	Monroe	14,932	21	20
Brown	2,154	7	7	Montgomery	6,402	<5	<5
Carroll	2,657	<5	<5	Morgan	11,334	26	23
Cass	6,910	5	5	Newton	2,330	<5	<5
Clark	17,945	8	8	Noble	7,542	19	19
Clay	4,431	<5	<5	Ohio	868	<5	<5
Clinton	6,565	<5	<5	Orange	3,239	<5	<5
Crawford	1,591	<5	<5	Owen	2,793	<5	<5
Daviess	4,901	7	7	Parke	2,309	<5	<5
Dearborn	8,682	25	25	Perry	3,014	<5	<5
Decatur	4,363	<5	<5	Pike	1,916	<5	<5
DeKalb	7,094	11	11	Porter	27,899	53	51
Delaware	16,237	10	10	Posey	3,695	<5	<5
DuBois	7,164	<5	<5	Pulaski	2,209	9	9
Elkhart	37,555	49	49	Putnam	5,876	<5	<5
Fayette	3,687	<5	<5	Randolph	5,684	<5	<5
Floyd	12,637	17	17	Ripley	5,613	5	5
Fountain	2,702	<5	<5	Rush	2,367	<5	<5
Franklin	2,516	<5	<5	Saint Joseph	40,862	29	27
Fulton	2,553	<5	<5	Scott	3,862	5	5
Gibson	5,169	<5	<5	Shelby	7,801	7	7
Grant	9,628	10	10	Spencer	3,272	7	7
Greene	5,083	6	6	Starke	3,732	11	11
Hamilton	62,159	53	52	Steuben	4,217	<5	<5
Hancock	14,443	16	16	Sullivan	3,294	<5	<5
Harrison	6,243	<5	<5	Switzerland	1,631	<5	<5
Hendricks	31,168	24	24	Tippecanoe	24,823	14	14
Henry	7,427	<5	<5	Tipton	2,449	<5	<5
Howard	14,583	15	15	Union	1,401	<5	<5
Huntington	5,340	5	5	Vanderburgh	23,896	15	15
Jackson	7,317	11	11	Vermillion	2,570	7	7
Jasper	5,228	7	6	Vigo	15,184	11	11
Jay	3,408	<5	<5	Wabash	5,790	<5	<5
Jefferson	4,507	9	9	Warren	1,377	<5	<5
Jennings	4,550	7	7	Warrick	10,610	<5	<5
Johnson	28,191	14	13	Washington	4,379	<5	<5
Knox	5,568	5	5	Wayne	11,023	5	5
Kosciusko	12,342	18	18	Wells	5,172	<5	<5
LaGrange	5,708	10	10	White	4,947	<5	<5
Lake	83,370	66	64	Whitley	6,375	8	8
LaPorte	17,745	13	12	Indiana	1,103,858	1,006	981
Lawrence	6,746	<5	<5				

Note: Incident numbers reflect each time a student was suspended/expelled due to alcohol use; unique count refers to the number of unique students involved (if the same student is suspended twice for alcohol, that reflects two incidents and one unique student).

Source: Indiana Department of Education, 2019

REFERENCES, CHAPTER 2

- Alcohol and Tobacco Commission. (2017). *Number of alcohol sales licenses in Indiana by county, 2016*. Data received January 10, 2019 from the Indiana Prevention Resource Center.
- Center for Behavioral Health Statistics and Quality (2016). *2015 National survey on drug use and health: Summary of the effects of the 2015 NSDUH questionnaire redesign: Implications for data users*. Substance Abuse and Mental Health Services Administration, Rockville, MD. Retrieved from <https://www.samhsa.gov/data/sites/default/files/NSDUH-TrendBreak-2015.pdf>
- Centers for Disease Control and Prevention. (1991-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (1999-2017). *CDC WONDER underlying causes of death (compressed mortality)*. Retrieved from <http://wonder.cdc.gov/>
- Centers for Disease Control and Prevention. (2006–2010). *Alcohol-related disease impact (ARDI)*. Retrieved from http://nccd.cdc.gov/DPH_ARDI/
- Centers for Disease Control and Prevention. (2016). *Fact sheets – Alcohol use and your health*. Retrieved from <https://www.cdc.gov/alcohol/fact-sheets/alcohol-use.htm>
- Centers for Disease Control and Prevention. (2018). *Behavioral Risk Factor Surveillance System (BRFSS) prevalence & trends data*. Retrieved from <http://www.cdc.gov/brfss/brfssprevalence/index.html>
- Gassman, R., Jun, M., Samuel, S., Agle, J. D., King, R., Ables, E., Wolf, J. (2018). *Indiana Youth Survey*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Indiana Department of Child Services. (2019). *SFY2018 Child Removals due to Parent Substance Abuse*. Source: MaGIK CHINS AFCARS; retrieved 2/1/2019.
- Indiana Department of Education. (2019). *DOE discipline 2018, incidents and student counts*. Aggregate data received from Jeff Wittman on January 18, 2019.
- Indiana Family and Social Services Administration. (2018). *Treatment Episode Data System (TEDS)*. Indianapolis, IN: Indiana Family and Social Services Administration.
- Indiana State Department of Health. (2015-2017). *Annual Legislative Report of the Indiana Birth Defects and Problems Registry, 2018*. Retrieved from https://www.in.gov/isdh/files/OPA_Approved_2018%20Annual%20Legislative%20Report%20for%20IBDPR.pdf
- Indiana State Department of Health. (2017). *Indiana hospital discharge data*. Retrieved from <http://www.in.gov/isdh/20624.htm>
- Indiana State Police. (2017). *Automated Reporting Information Exchange System (ARIES), Vehicle Crash Records System*. Data received from the Center for Criminal Justice Research, Public Policy Institute, School of Public and Environmental Affairs, Indiana University–Purdue University Indianapolis.
- Inter-university Consortium for Political and Social Research, University of Michigan. (2018). *Monitoring the Future (MTF)*. Retrieved from <http://www.monitoringthefuture.org/data/data.html>
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey, 2018*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey>
- National Institute on Alcohol Abuse and Alcoholism. (2018). *Apparent per capita alcohol consumption: National, state, and regional trends, 1977–2016*. Retrieved from <https://www.openicpsr.org/openicpsr/project/105583/version/V1/view>
- Sacks, J. J., Gonzales, K. R., Bouchery, E. E., Tomedi, L. E., & Brewer, R. D. (2015). 2010 National and state costs of excessive alcohol consumption. *American Journal of Preventive Medicine*, 49(5), e73-e79.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2016). *Treatment Episode Data Set -- Admissions (TEDS-A)*. Retrieved from <http://www.dasis.samhsa.gov/dasis2/teds.htm>

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>

Substance Abuse and Mental Health Services Administration, Fetal Alcohol Spectrum Disorders Center for Excellence. (2007). *Fetal Alcohol Spectrum Disorders by the numbers*. Retrieved from http://www.fascenter.samhsa.gov/documents/WYNK_Numbers.pdf

TOBACCO USE IN INDIANA: CONSUMPTION PATTERNS AND CONSEQUENCES

INTRODUCTION

In the United States, one of every five deaths is related to cigarette smoking, making it the leading cause of preventable disease and death (U.S. Department of Health and Human Services [USDHHS], 2014). The adverse effects of tobacco on population health have been well-researched. In Indiana, more than 11,100 adults die every year from their own smoking, and 333,000 live with a tobacco-related disease (USDHHS, 2014). Furthermore, 151,000 (approximately 1 in 10) Indiana youth now under the age of 18 will die prematurely from a smoking-related illness (USDHHS, 2014). Additionally, over 1,300 adults, children, and infants died in 2014 as a result of exposure to secondhand smoke (Lewis & Zollinger, 2014). Indiana incurs close to \$3 billion annually in healthcare costs directly caused by smoking, including nearly \$590 million that is absorbed by Medicaid (Campaign for Tobacco-Free Kids, 2018b).

Though self-reported cigarette smoking has been on the decline, electronic nicotine delivery systems, including e-cigarettes, have surged in popularity in recent years


(Marynak et al., 2017). While e-cigarettes have been promoted as less dangerous than cigarettes, they have not been approved as safe by the U.S. Food and Drug Administration (FDA) and long-term health effects of exposure to aerosol from e-cigarettes are currently unknown (Indiana State Department of Health, Tobacco Prevention and Cessation Commission [ISDH/TPCC], 2018a).

PREVALENCE OF TOBACCO CONSUMPTION IN THE GENERAL POPULATION

National Survey on Drug Use and Health

Estimates from the 2017 National Survey on Drug Use and Health (NSDUH) showed that 26.8% (95% Confidence Interval [CI]: 24.6 - 29.1) of Indiana residents 12 years and older used a tobacco product in the past month, a rate significantly higher than the U.S. rate (23.0%; 95% CI: 22.6–23.4). Tobacco products include cigarettes, smokeless tobacco, cigars, and pipe tobacco. Indiana’s rate has gradually decreased over the past decade (see Figure 3.1) (Substance Abuse and Mental Health Services Administration [SAMHSA], 2017).

Figure 3.1 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Any Tobacco Use in the Past Month (National Survey on Drug Use and Health, 2007–2017)




Source: SAMHSA, 2017

Among tobacco users, the most commonly used type of tobacco was cigarettes. In 2017, 21.4% (95% CI: 19.3-23.6) of Hoosiers ages 12 years and older reported past-month use of cigarettes, a rate


significantly higher than the U.S. rate (18.5%; 95% CI: 18.1–18.8). Indiana’s smoking prevalence declined from 28.0% in 2007 (95% CI: 25.6–30.4) to 21.4% in 2017 (95% CI: 19.3-23.6) (see Figure 3.2).

Figure 3.2 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Cigarette Use in the Past Month (National Survey on Drug Use and Health, 2007–2017)



Source: SAMHSA, 2017

Figure 3.3 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Cigarette Use in the Past Month (National Survey on Drug Use and Health, 2017)



Source: SAMHSA, 2017

Tobacco use differed by age group and was most prevalent among young adults. Over 28 percent of 18- to 25-year-olds in Indiana reported smoking cigarettes in the past month (95% CI: 20.9-27.9) compared to 24.3% of their national same-age counterparts (95% CI: 22.3-23.6) (see Figure 3.3) (SAMHSA, 2017).

Behavior Risk Factor Surveillance System

The Behavioral Risk Factor Surveillance System (BRFSS) focuses on behaviors and conditions that are linked to the leading causes of death. According to 2017 findings, the prevalence rate for adult smoking in Indiana was 21.8% (95% CI: 20.8–22.8). Moreover, 15.8% (95% CI: 14.9–16.7) of Hoosiers reported using cigarettes every day. Indiana’s smoking rates were higher than the national median rates; i.e., 17.0% of U.S. adults smoked in the past month and 11.7% reported smoking every day (Centers for Disease Control and Prevention [CDC], 2018a). Statistically significant differences in smoking prevalence were observed for the following groups in Indiana (see Table 3.1):

- Smoking rates were higher among men than women.
- Smoking was less prevalent in Hispanic Hoosiers compared to those who identified as white or black.
- Smoking prevalence was lowest among older adults ages 65 and above.
- Educational attainment was inversely associated with prevalence rate, i.e., individuals who achieved higher levels of education had lower smoking rates.
- Income level was inversely associated with prevalence rate, i.e., individuals with higher income levels had lower smoking rates.

Adult smoking prevalence in Indiana continues to be above the U.S. level (see Figure 3.4).

Table 3.1 Adult Smoking Prevalence in Indiana, by Gender, Race/Ethnicity, Age Group, Educational Attainment, and Income Level (Behavioral Risk Factor Surveillance System, 2017)

		Indiana (95% CI)
Gender	Male	23.5% (22.0 - 25.0)
	Female	20.2% (18.8 - 21.5)
Race / Ethnicity	White	22.3% (21.2 - 23.5)
	Black	21.3% (17.7 - 25.0)
	Hispanic	14.0% (10.3 - 17.6)
Age Group	18-24	18.7% (15.1 - 22.3)
	25-34	28.4% (25.3 - 31.5)
	35-44	26.1% (23.4 - 28.9)
	45-54	26.0% (23.7 - 28.4)
	55-64	23.3% (21.4 - 25.2)
	65+	10.4% (9.3 - 11.5)
Education	Less than High School	36.8% (32.8 - 40.9)
	High School or GED	26.5% (24.7 - 28.3)
	Some post-High School	21.1% (19.3 - 22.8)
	College Graduate	7.5% (6.5 - 8.5)
Income	Less than \$15,000	35.7% (31.9 - 39.4)
	\$15,000–\$24,999	32.5% (29.7 - 35.4)
	\$25,000–\$34,999	25.6% (22.3 - 28.9)
	\$35,000–\$49,999	22.7% (19.8 - 25.6)
	\$50,000 and above	14.2% (12.9 - 15.5)
Total		21.8% (20.8 - 22.8)

Source: CDC, 2017a

Indiana Adult Tobacco Survey

The 2017 Indiana Adult Tobacco Survey (IATS) estimated the overall smoking prevalence among Indiana adults at 23.6% (95% CI: 21.1–26.2). Smoking was most prevalent among persons:


- Without a high school degree (49.0%; 95% CI: 39.6–58.5)
- With annual household incomes less than \$20,000 (42.4%; 95% CI: 32.7–52.7)
- Ages 25 to 39 years (32.9%; 95% CI: 27.0–39.4)

- Who are non-Hispanic black (30.9%; 95% CI: 22.5–40.8)

Approximately 23.7% (95% CI: 21.2–26.3) of adults in Indiana reported ever trying an e-cigarette.

Among current smokers, nearly one-third (31.5%; 95% CI: 25.7–38.0) reported intentions to quit within the next 30 days (Indiana State Department of Health [ISDH], Tobacco Prevention & Cessation Commission [TPCC], 2019). For details on smokers' intentions to quit, see Table 3.2.

Figure 3.4 Percentage of Indiana and U.S. Population (18 Years and Older) Reporting Current Cigarette Use (Behavioral Risk Factor Surveillance System, 2011–2017)



Source: CDC, 2017a

Table 3.2 Intentions to Quit Smoking Among Current Smokers (Indiana Adult Tobacco Survey, 2017)

	Within next 30 days	Within 30 days to 6 months	Sometime after 6 months	No intention to quit
Gender				
Male	28.2% (20.4–37.6)	15.3% (9.7–23.4)	14.6% (9.1–22.7)	41.9% (32.9–51.4)
Female	35.7% (27.3–45.0)	16.4% (10.8–24.0)	20.3% (14.0–28.5)	27.7% (20.3–36.4)
Race/Ethnicity				
White	28.4% (22.1–35.7)	15.3% (10.7–21.5)	17.7% (12.7–24.1)	38.6% (31.6–46.1)
Black	48.8% (31.8–66.1)	19.3% (8.9–36.8)	22.2% (10.7–40.4)	9.8% (4.0–21.8)
Hispanic	28.5% (7.2–67.2)	12.1% (1.6–53.3)	10.6% (1.4–49.6)	48.7% (17.9–80.6)
Other	43.0% (17.7–72.5)	10.8% (2.4–37.1)	20.6% (5.4–54.2)	25.7% (8.2–57.1)
Age Group				
18-24	23.0% (12.5–38.4)	14.3% (6.4–28.8)	17.7% (8.1–34.4)	45.0% (30.0–61.0)
25-39	34.2% (23.5–46.8)	17.8% (10.3–29.1)	18.4% (10.8–29.4)	29.6% (19.9–41.5)
40-64	33.6% (24.9–43.5)	14.4% (9.1–22.1)	18.8% (12.4–27.4)	33.2% (24.6–43.2)
65+	29.7% (15.8–48.7)	17.8% (7.1–37.9)	12.7% (3.7–35.3)	39.9% (23.2–59.4)
Education				
Less than High School	35.9% (22.3–52.1)	18.7% (9.2–34.5)	11.2% (5.0–23.3)	34.2% (21.2–50.1)
High School Grad	26.1% (18.3–35.8)	14.9% (9.1–23.4)	23.9% (15.9–34.4)	35.0% (25.8–45.5)
Some College	34.8% (24.6–46.6)	12.7% (7.2–21.3)	18.4% (11.1–28.9)	34.1% (24.5–45.3)
College	32.4% (16.6–53.5)	25.2% (12.0–45.4)	10.5% (2.6–33.6)	31.9% (15.9–53.7)
Post-Graduate	27.7% (8.3–61.9)	10.0% (1.9–39.0)	2.5% (0.5–10.7)	59.8% (28.8–84.6)
Income				
Less than \$20,000	39.0% (23.6–57.0)	13.3% (5.6–28.1)	18.6% (8.0–37.4)	29.2% (16.9–45.6)
\$20,000 – \$39,999	33.0% (22.1–46.0)	17.4% (9.4–30.0)	25.8% (16.6–37.8)	23.8% (15.3–35.1)
\$40,000 – \$69,999	32.5% (22.0–45.0)	18.8% (11.2–29.9)	14.6% (7.8–25.7)	34.2% (23.3–46.9)
\$70,000 or more	26.1% (17.1–37.8)	12.5% (6.9–21.5)	13.1% (6.9–23.4)	48.3% (36.6–60.2)
Total	31.5% (25.7–38.0)	15.7% (11.5–21.0)	17.9% (13.4–23.5)	34.9% (28.9–41.4)

Source: ISDH/TPCC, 2019


Indiana Youth Tobacco Survey

The Indiana Youth Tobacco Survey (IYTS) is a statewide school-based survey of middle school (grades 6 through 8) and high school (grades 9 through 12) students that captures information on various tobacco-related issues, such as tobacco use, smoking cessation, tobacco-related attitudes and beliefs, social influences on tobacco use, and secondhand smoke exposure. According to IYTS results, cigarette, smokeless tobacco products, and

overall tobacco use declined significantly in Indiana from 2004 to 2016 (see Figures 3.5 and 3.6) (ISDH/TPCC, 2019).

Based on 2016 IYTS results, a total of 4.9% of middle school students (95% CI: 3.5–6.0) and 20.3% of high school students (95% CI: 15.4–25.6) used any tobacco product in the past month. Among middle school students, 1.8% (95% CI: 1.9–3.8) and among high school students, 8.7% (95% CI: 8.6–15.4) reported smoking cigarettes in


Figure 3.5 Tobacco Use among Indiana High School Students (9th–12th Grade) (Indiana Youth Tobacco Survey, 2004–2016)



Note: Due to the emergence of new tobacco products in recent years and corresponding changes to the survey instrument, the definition of “any tobacco use” has changed over time. Between 2004 and 2010, “any tobacco use” included cigarettes, cigars, smokeless tobacco, pipe, or bidis. In 2012, 2014, and 2016 “any tobacco use” included cigarettes, cigars, smokeless tobacco, bidis, pipe, hookah, snus, dissolvable tobacco, and e-cigarettes.

Source: ISDH/TPCC, 2019

Figure 3.6 Percentage of Indiana Middle School and High School Students Reporting Current Tobacco and Cigarette Use (Indiana Youth Tobacco Survey, 2004–2016)



Source: ISDH/TPCC, 2018

the past month. In 2016, 2.8% of middle school students and 10.5% of high school students in Indiana reported current use of e-cigarettes. Among Indiana youth who currently smoke cigarettes, 33.6% of middle school students and 45.8% of high school students also reported currently using e-cigarettes (ISDH/TPCC, 2019).

Appendix 3A shows the percentages, including 95% confidence intervals, of Indiana middle and high school students who reported current use of various tobacco products, grouped by gender, race/ethnicity, and grade, in 2016.

Youth Risk Behavior Surveillance System

The use of tobacco products has wide-ranging consequences for adolescents and young adults. Factors associated with youth tobacco use include low socioeconomic status; use and approval of tobacco use by peers or siblings; smoking by parents or guardians; accessibility, availability and price of tobacco products; a perception that tobacco use is normative; lack of parental support or involvement; low levels of academic achievement; lack of skills to resist influences to tobacco use; lower self-image or self-esteem; belief in functional benefits of tobacco use; and lack of self-efficacy to refuse offers of tobacco (CDC, 2016b).

The Youth Risk Behavior Surveillance System (YRBSS) monitors health-risk behaviors such as


tobacco, alcohol, and other drug use, which contribute to death and disability among youths in schools nationwide. According to 2015 YRBSS findings, almost one-third of high school students currently use a tobacco product, primarily electronic vapor products (vapor products) (see Table 3.3). In Indiana, rates of current cigarette use decreased significantly from 25.6% (95% CI: 23.2–28.2) in 2003 to 11.2% (95% CI: 8.3–14.8) in 2015; however, vapor products have gained popularity with nearly one-fourth of high school students (23.9%; 95% CI: 20.6–27.7) reporting current use (CDC, 1991-2015). For more information, see Figures 3.7 through 3.9.

Table 3.3 Current Use of Tobacco Products in Indiana and U.S. High School Students (Youth Risk Behavior Surveillance System, 2015)

	Indiana (95% CI)	U.S. (95% CI)
Any Tobacco Use	32.4% (27.3–38.0)	31.4% (29.1–33.8)
Electronic Vapor Products	23.9% (20.6–27.7)	24.1% (22.1–26.2)
Cigarettes	11.2% (8.3–14.8)	10.8% (9.4–12.4)
Cigars	11.4% (9.1–14.3)	10.3% (9.0–11.8)
Smokeless Tobacco	9.4% (5.9–14.7)	7.3% (6.1–8.6)


Source: CDC, 1991-2015

Figure 3.7 Rates of Current Use of Cigarettes and Electronic Vapor Products in Indiana High School Students (9th–12th Grade), by Gender (Youth Risk Behavior Surveillance System, 2015)




Source: CDC, 1991-2015

Figure 3.8 Rates of Current Use of Cigarettes and Electronic Vapor Products in Indiana High School Students (9th–12th Grade), by Race/Ethnicity (Youth Risk Behavior Surveillance System, 2015)



Source: CDC, 1991-2015

Figure 3.9 Rates of Current Use of Cigarettes and Electronic Vapor Products in Indiana High School Students (9th -12th Grade), by Grade (Youth Risk Behavior Surveillance System, 2015)




Source: CDC, 1991-2015

Indiana Youth Survey

The Indiana Youth Survey, conducted annually of students in grades 6 to 12, assesses students' substance use, mental health, gambling, and risk and protective factors that can affect their academic success. Findings from the 2018 survey showed that tobacco use increased as students progressed in school, i.e., higher smoking

rates occurred among 12th grade students than 8th graders, both for cigarettes and electronic vapor products (such as e-cigarettes, vaping pens, and e-hookahs) (see Figure 3.10) (Gassman et al., 2018). See Appendix 3B for Indiana students' 2018 monthly cigarette and vaping products use by region and grade.

Figure 3.10 Monthly Cigarette Use and Vaping among 8th, 10th, and 12th Grade Students, Indiana and the United States (Indiana Youth Survey and Monitoring the Future Survey, 2018)




Source: Gassman et al., 2018; Inter-university Consortium for Political and Social Research, 2018

Comparisons between Indiana and the United States on 30-day prevalence of cigarette use and vaping among 12th grade students imply that (a) Hoosier students have had higher rates throughout the years, and (b) cigarette use has been declining, while vaping rates are

at an all-time high (see Figure 3.11). However, these results need to be interpreted with caution, as statistical significance could not be determined due to the lack of detail provided in the publicly available data sets.

Figure 3.11 Monthly Cigarette Use and Vaping among 12th Grade Students in Indiana and the United States (Indiana Youth Survey: 2008–2018; and Monitoring the Future Survey, 2008–2018)



Note: Vaping data only available since 2015.

Source: Gassman et al., 2018; Inter-university Consortium for Political and Social Research, 2018

Indiana College Substance Use Survey

The Indiana College Substance Use Survey includes questions on the use of various tobacco products. The 2018 survey, which was based on 24 participating colleges and universities, showed that electronic vapor products were the most commonly used nicotine delivery system, with 21.8% of Indiana college students reporting current (past-month) use (U.S.: 11.3%); followed by cigarettes, the second most common form (Indiana: 12.4%; U.S.: 8.0%). Consumption rates for the different types of tobacco/nicotine products by demographic characteristics can be found in Table 3.4 (King & Jun, 2018).¹

CONSEQUENCES OF TOBACCO USE

The use of tobacco can lead to tobacco/nicotine dependence as well as tobacco-related diseases (CDC, 2017b). The risk of developing serious health problems associated with tobacco significantly decreases as people quit using tobacco products. Several factors influence tobacco cessation including healthcare coverage/costs, socioeconomic characteristics, availability of tobacco cessation products and media campaigns.

Additionally, tobacco use in K-12 students on school property or during school activities can lead to disciplinary actions, including suspensions and expulsions. During academic year 2018, a total of 4,817 suspensions/expulsions were recorded in Indiana schools involving tobacco use (Indiana Department of Education, 2019). For the number of tobacco-related incidents by county, see Appendix 3C.

Tobacco-Related Morbidity

Smoking affects respiratory health and is related to chronic coughing and wheezing among adults. Smokers are more likely than nonsmokers to have upper and lower respiratory tract infections. Generally, lung function deteriorates more quickly in smokers than in nonsmokers. Smoking contributes significantly to the number of deaths from lung cancer, heart disease, chronic lung diseases, and other illnesses (USDHHS, 2014). Adverse outcomes of smoking also include cancers of the oral cavity, pharynx, larynx, esophagus, bladder, stomach, cervix, kidney, and pancreas. Furthermore, smoking has been linked to liver, colorectal, prostate, and breast cancers, and can also result in acute myeloid leukemia (USDHHS, 2014). For smoking-attributable cancers, the risk generally increases with the number of cigarettes smoked and the number of years of smoking, and usually decreases after the smoker quits completely. The leading cause of cancer deaths is lung cancer, and cigarette smoking causes most cases. However, any tobacco use can be detrimental. Smokeless tobacco has been shown to cause oral cancers and may also be a risk factor for cardiovascular disease (CDC, 2016a). Other specific health-related outcomes include age-related macular degeneration, dental disease, diabetes, autoimmune disease, rheumatoid arthritis, systemic lupus erythematosus, and inflammatory bowel disease (USDHHS, 2014).

Smoking may harm men's and women's reproductive health, and the effects can be seen in fetuses, infants, and children. Smoking can affect men's sperm and lead to reduced fertility and increased risk for

Table 3.4 Rates of Past-Month Use of Nicotine Products among Indiana College Students (Indiana College Substance Use Survey, 2018)


	Indiana (Total)	Male	Female	Under 21	21 or Over
Cigarettes	12.4	15.6	10.3*	10.8	15.1*
Cigars	6.8	12.3	3.6*	6.7	7.0
Chewing/smokeless tobacco	3.5	8.0	0.8*	3.3	3.9
Smoking tobacco with hookah/ water pipe	5.7	6.6	5.0*	5.0	6.8*
Electronic vapor products	21.8	28.2	17.9*	24.2	17.8

Note: * $P < 0.05$

Source: King & Jun, 2018

¹Twenty-four Indiana colleges participated in the 2018 survey; results are based on nonrandom sampling and are not representative of all college students in Indiana.

Figure 3.12 Percentage of Smoke-free Homes and Workplaces in Indiana (Adult Tobacco Survey, 2002–2017)



Source: ISDH/TPCC, 2019

birth defects and miscarriage. Women who smoke have an increased risk for infertility and ectopic pregnancies. Smoking during pregnancy results in health problems for both mothers and babies. These include increased risk of spontaneous abortions, pregnancy complications (e.g., placenta previa, placental abruption, and premature rupture of membranes before labor begins), premature delivery, low birth-weight infants, stillbirth, and sudden infant death syndrome (SIDS). Mothers who smoke during pregnancy reduce their babies' lung function (CDC, 2016a). In Indiana, the percentage of births to mothers who smoked during pregnancy declined from 18.5% in 2007 to 13.5% in 2017; a higher percentage of white mothers (14.8%) smoked during pregnancy than black mothers (10.1%) in 2017 (ISDH/Epidemiology Resource Center, 2018). The Indiana State Department of Health, Tobacco Prevention and Cessation provides county-level information on various smoking-related outcomes. For a detailed list, see Appendix 3D.

Secondhand smoke: Secondhand smoke (sometimes called environmental tobacco smoke) has serious health consequences. An estimated 58 million nonsmoking Americans continue to be exposed to secondhand smoke in homes, vehicles, workplaces, and public places. Exposure to tobacco smoke can cause heart disease and lung cancer even in nonsmoking adults, increasing the risk by 25% to 30% for heart disease and by 20% to 30% for lung cancer. Children, in particular, are heavily impacted by secondhand

smoke. Exposure increases their chance of developing significant lung conditions, especially asthma and bronchitis. Also, secondhand smoke can cause SIDS, acute respiratory infections, ear problems, and more frequent and severe asthma attacks in children. In the United States, secondhand smoke is responsible for nearly 34,000 deaths due to heart disease, more than 8,000 deaths from stroke, and over 7,300 lung cancer deaths each year among nonsmoking adults (USDHHS, 2014). An estimated 1,337 Hoosiers die each year from secondhand smoke (Lewis & Zollinger, 2014).

In Indiana, the percentage of smoke-free homes has increased from 60.1% in 2002 to 78.2% in 2017. The percentage of smoke-free workplaces² rose from 60.3% to 94.6% during that time period (see Figure 3.12). Although Indiana is making progress, it is lagging behind the rest of the nation terms of *comprehensive* coverage from secondhand smoke exposure (comprehensive coverage includes workplaces, restaurants, and bars). With the addition of the statewide smoke-free air law in 2012, all Indiana residents are covered in most workplaces and restaurants, but the law exempts bars, clubs, and gaming facilities. As of January 2019, a total of 23 communities³ in Indiana have passed comprehensive smoke-free air ordinances which cover all workplaces, including bars, to ensure that all workers are protected from secondhand smoke. These 23 comprehensive ordinances cover approximately 31% of all residents in Indiana (ISDH/TPCC, 2019).

²This measure refers to the prevalence of workers reporting a 100% smoke-free workplace (Adult Tobacco Survey).

³These are Delaware Co., Hancock Co., Howard Co., Monroe Co., Vanderburgh Co., Vigo Co., Austin, Bloomington, Columbus, Cumberland, Elkhart, Fort Wayne, Franklin, Greencastle, Indianapolis, Kokomo, Lawrence, North Manchester, Plainfield, South Bend, Terre Haute, West Lafayette, and Zionsville.

E-cigarettes: Research shows that e-cigarette aerosol releases measurable amounts of carcinogens and other toxins into the air, including nicotine, formaldehyde, and acetaldehyde. In addition, e-cigarette aerosol has been found to contain a high concentration of ultra-fine particles. Exposure to fine and ultra-fine particles may exacerbate respiratory conditions and constrict arteries. In addition, nicotine from e-cigarettes may lead to increased heart rate and diastolic blood pressure. (ISDH/TPCC, 2018a).

E-cigarettes are the most commonly used tobacco product among youth in Indiana and nationwide. There is substantial evidence that e-cigarette use increases the risk of using regular combustible cigarettes among youth and young adults. For example, nearly half of Indiana high school students who used e-cigarettes in 2016 also smoked regular cigarettes, and the percentage of Hoosier adults reporting dual use was 58.5% (ISDH/TPCC, 2018a).

In 2016, the U.S. Surgeon General issued a report highlighting concerns related to vaping among youth and young adults (USDHHS, 2016). Key findings of the report are as follows:

- E-cigarette use among youth and young adults has become a public health concern.
- E-cigarettes are the most commonly used tobacco product among youth, and use of e-cigarettes is strongly associated with use of other tobacco products.
- The use of products containing nicotine pose danger to youth, pregnant women, and fetuses. The use of products containing nicotine among youth, including e-cigarettes, is unsafe.
- E-cigarette aerosol is not harmless. It can contain harmful and potentially harmful constituents.
- E-cigarettes are marketed by promoting flavors and using a variety of media channels and approaches that have been used in the past to market tobacco to youth and young adults.

A new group of e-cigarette products look like USB drives. The most popular brand, JUUL (pronounced “jewel”), has grown quickly in popularity since introduction to the market in 2015, fueled by a large following among youth and young adults. Because of its unsuspecting appearance and small size, JUUL devices may not be immediately identified as an e-cigarette,

and can be easily concealed. The increased use of these products has become a concern for educators in Indiana. Many report that students are concealing JUUL and using it in schools. Nicotine use can have adverse effects on adolescent brain development. Therefore, nicotine use by youth in any form is unsafe, and efforts are warranted to educate youth about the dangers of use of all forms of tobacco products, regardless of whether they are combustible, noncombustible, or electronic. The skyrocketing e-cigarette use rate among youth observed in the past year has been partially attributed to the surge in JUUL popularity. The Surgeon General issued an advisory in December 2018 stressing the importance of protecting children from a lifetime of nicotine addiction and associated health risks by immediately addressing the epidemic of youth e-cigarette use.

Tobacco-Related Mortality

As the second major cause of death in the world, tobacco is responsible for approximately 6 million deaths every year, including about 600,000 deaths from exposure to secondhand smoke (World Health Organization, 2015). In the United States, cigarette smoking is the single most preventable cause of disease and death, causing more deaths annually than acquired immune deficiency syndrome (AIDS), alcohol, cocaine, heroin, homicide, suicide, motor vehicle crashes, and fires combined (USDHHS, 2014).

In the United States, tobacco use is responsible for more than 480,000 deaths per year among adults age 35 and older. In addition, 16 million adults are suffering from smoking-related conditions. On average, smoking reduces adult life expectancy by a minimum of 10 years. Smoking is the leading risk-factor for lung cancer, which is the foremost cause of cancer-related deaths for both males and females (Siegel, Miller, & Jemal, 2015).

Economic Impact

In 2016, the annual U.S. tobacco industry marketing expenditures were approximately \$9.5 billion, including Indiana’s share of \$296.8 million. The state’s total tobacco marketing expenditures declined after peaking at \$475.4 million in 2003 (Campaign for Tobacco-Free Kids, 2018b; Federal Trade Commission, 2018).

The federal excise tax is \$1.01 per pack of cigarettes. The average state cigarette excise tax is \$1.79 per pack, but varies from 17 cents in Missouri to

\$4.50 in Washington DC; Indiana's tobacco excise tax rate is 99.5 cents per pack (Campaign for Tobacco-Free Kids, 2018a).

Cigarette smoking is estimated to be responsible for greater than \$300 billion in annual health-related economic losses in the United States (\$170 billion in direct medical costs and approximately \$156 billion in lost productivity) (CDC, 2016a). In Indiana, \$2.93 billion dollars of health-related costs in 2009 were smoking-attributable expenditures (SAE). Most of these

costs accrued through hospital care (\$1.57 billion) and prescription drugs (\$525 million); the SAE estimate also included ambulatory care (\$405 million), nursing home care (\$283 million), and other health-related costs (\$147 million) (CDC, 2016a). The combination of increased medical costs, higher insurance rates, added maintenance expenses, lower productivity, and higher rates of absenteeism due to smoking adds financial strain to U.S. businesses every year.

APPENDIX 3A

Percentage of Indiana Middle School and High School Students Who Currently Use Cigarettes, E-Cigarettes, or Smokeless Tobacco by Gender, Race/Ethnicity, and School Grade (Indiana Youth Tobacco Survey, 2016)

	Current Use of Cigarettes		Current Use of E-Cigarettes		Current Use of Smokeless Tobacco	
	%	(95% CI)	%	(95% CI)	%	(95% CI)
MIDDLE SCHOOL						
Gender						
Male	1.4	(0.8–2.1)	2.3	(1.4–3.2)	1.7	(0.7–2.7)
Female	2.1	(0.8–3.4)	3.3	(2.2–4.3)	0.8*	(0.1–1.6)
Race/Ethnicity						
White	1.8	(0.6–3.0)	2.9	(1.7–4.0)	1.5	(0.8–2.2)
Black	1.2*	(0.0–2.4)	2.4*	(0.0–5.0)	0.7*	(0.0–2.0)
Hispanic	2.8	(1.4–4.1)	4.2	(2.1–6.4)	1.2*	(0.0–2.3)
Grade						
6	0	(0.0–0.0)	1.5*	(0.2–2.9)	1.5*	(0.3–2.7)
7	1.3	(0.5–2.1)	1.7*	(0.6–2.9)	0.9*	(0.2–1.7)
8	3.2	(1.2–5.3)	4.5	(2.3–6.7)	1.4	(0.1–2.7)
Total	1.8	(1.0–2.5)	2.8	(1.9–3.7)	1.3	(0.7–1.9)
HIGH SCHOOL						
Gender						
Male	9.3	(6.8–11.9)	12.0	(8.6–15.3)	8.1	(5.0–11.1)
Female	8.2	(5.7–10.8)	9.1	(7.3–10.9)	2.3	(0.9–3.7)
Race/Ethnicity						
White	10.0	(7.7–12.3)	11.6	(9.2–14.0)	6.1	(3.8–8.4)
Black	3.1*	(0.4–5.8)	3.9	(1.9–5.9)	1.2*	(0.0–3.0)
Hispanic	8.0	(4.9–11.1)	11.7	(7.4–15.9)	2.8*	(0.8–4.7)
Grade						
9	4.4	(2.3–6.4)	6.2	(3.0–9.4)	3.3	(1.5–5.1)
10	8.1	(5.5–10.7)	10.8	(7.9–13.7)	4.8	(2.4–7.3)
11	9.7	(5.9–13.5)	9.5	(6.4–12.6)	6.6*	(2.5–10.7)
12	13.0	(8.3–17.6)	15.8	(11.2–20.3)	6.2	(3.2–9.2)
Total	8.7	(6.7–10.8)	10.5	(8.4–12.5)	5.3	(3.3–7.2)

Note: *Indicates data are statistically unstable because the relative standard error is >30%. These estimates should be interpreted with caution.

Source: ISDH/TPCC, 2019

APPENDIX 3B - Part 1

Percentage of Indiana Students Reporting Monthly Cigarette Use, by Region and Grade (Indiana Youth Survey, 2018)

	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
6th Grade	1.2	0.7*	1.2	1.1	1.7	1.2	1.3	1.1	1.5
7th Grade	2.4	1.5*	2.5	1.9	2.6	2.3	2.6	1.9	3.3*
8th Grade	4.0	2.7*	4.0	5.3	3.5	3.5	5.3*	3.4	5.2*
9th Grade	5.1	4.4	4.4	4.8	4.9	3.6*	5.7	5.8	6.3*
10th Grade	6.8	5.7*	5.6*	7.2	5.9	7.7	7.5	8.5*	7.5
11th Grade	8.6	6.8*	5.7*	6.7	8.3	8.0	10.4*	11.1*	10.1*
12th Grade	9.9	8.8	7.9*	8.6	9.7	8.8	9.0	11.7*	12.1*

Note: * Indicates a local rate that is significantly different from the overall state rate ($P < 0.05$).

Source: Gassman et al., 2018

APPENDIX 3B - Part 2

Percentage of Indiana Students Reporting Monthly E-Cigarette Use, by Region and Grade (Indiana Youth Survey, 2018)

	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
6th Grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
7th Grade	7.0	6.9	7.6	3.7*	6.4	7.7	5.4*	7.0	8.1*
8th Grade	11.9	14.1*	11.3	10.5	9.9*	12.6	10.3*	12.0	12.1
9th Grade	16.7	19.2*	15.3	12.6*	14.3*	14.2*	15.4	19.9*	18.1
10th Grade	20.4	22.4*	16.8*	15.1*	19.8	21.1	17.3*	25.9*	20.5
11th Grade	23.7	22.1	20.2*	18.1*	21.5	24.3	21.6	31.7*	24.9
12th Grade	28.6	28.7	23.6*	16.6*	24.0*	27.6	26.5	36.4*	32.4*

Note: * Indicates a local rate that is significantly different from the overall state rate ($P < 0.05$).

The Indiana Youth Survey did not ask 6th grade students about e-cigarette use.

Source: Gassman et al., 2018

APPENDIX 3C

Number of Incidents and Unique Students Involved in Suspensions/Expulsions due to Tobacco Use in Indiana,
Academic Year 2018

County	Students Enrolled	Number of Incidents	Number of Unique Students Involved
Adams	4,347	<5	<5
Allen	57,046	171	163
Bartholomew	13,126	135	125
Benton	1,928	14	14
Blackford	1,764	20	18
Boone	12,342	55	55
Brown	2,154	12	9
Carroll	2,657	14	14
Cass	6,910	41	40
Clark	17,945	69	66
Clay	4,431	<5	<5
Clinton	6,565	16	16
Crawford	1,591	36	34
Daviess	4,901	7	7
Dearborn	8,682	104	92
Decatur	4,363	8	8
DeKalb	7,094	49	47
Delaware	16,237	49	49
Dubois	7,164	41	40
Elkhart	37,555	136	130
Fayette	3,687	24	23
Floyd	12,637	162	156
Fountain	2,702	<5	<5
Franklin	2,516	19	18
Fulton	2,553	16	16
Gibson	5,169	11	11
Grant	9,628	43	40
Greene	5,083	32	29
Hamilton	62,159	415	391
Hancock	14,443	67	62
Harrison	6,243	72	61
Hendricks	31,168	185	169
Henry	7,427	59	53
Howard	14,583	40	36
Huntington	5,340	59	53
Jackson	7,317	31	31
Jasper	5,228	20	18
Jay	3,408	37	35
Jefferson	4,507	59	53
Jennings	4,550	<5	<5
Johnson	28,191	127	118
Knox	5,568	53	45
Kosciusko	12,342	147	133
LaGrange	5,708	23	23
Lake	83,370	267	254
LaPorte	17,745	66	63

(Continued on next page)

APPENDIX 3C (Continued from previous page)

County	Students Enrolled	Number of Incidents	Number of Unique Students Involved
Lawrence	6,746	56	47
Madison	20,089	128	116
Marion	179,578	342	327
Marshall	7,759	16	15
Martin	1,443	<5	<5
Miami	7,480	38	36
Monroe	14,932	92	84
Montgomery	6,402	55	50
Morgan	11,334	37	36
Newton	2,330	18	14
Noble	7,542	58	51
Ohio	868	<5	<5
Orange	3,239	9	9
Owen	2,793	40	33
Parke	2,309	<5	<5
Perry	3,014	6	5
Pike	1,916	14	14
Porter	27,899	122	110
Posey	3,695	23	22
Pulaski	2,209	34	31
Putnam	5,876	17	16
Randolph	5,684	18	17
Ripley	5,613	66	60
Rush	2,367	10	10
Scott	40,862	67	65
Shelby	3,862	15	13
Spencer	7,801	24	24
St. Joseph	3,272	<5	<5
Starke	3,732	25	23
Steuben	4,217	19	19
Sullivan	3,294	13	13
Switzerland	1,631	<5	<5
Tippecanoe	24,823	66	64
Tipton	2,449	14	13
Union	1,401	<5	<5
Vanderburgh	23,896	48	42
Vermillion	2,570	7	7
Vigo	15,184	6	6
Wabash	5,790	36	33
Warren	1,377	<5	<5
Warrick	10,610	59	56
Washington	4,379	59	55
Wayne	11,023	48	45
Wells	5,172	33	29
White	4,947	20	20
Whitley	6,375	48	47
Indiana	1,103,858	4,817	4,495

Note: Incident numbers reflect each time a student was suspended/expelled due to tobacco use; unique count refers to the number of unique students involved (if the same student is suspended twice for tobacco, that reflects two incidents and one unique student).

Source: Indiana Department of Education, 2019

APPENDIX 3D - Part 1

Adult Smoking Prevalence and Chronic Disease Outcomes, by County

County	Estimated adult smoking rate (Statewide: 2017 BRFSS; County-level: 2013-2017 BRFSS)	Age-adjusted rate of lung cancer deaths per 100,000 population (2013-2017)	Age-adjusted rate of major cardiovascular diseases deaths per 100,000 population (2013-2017)	Asthma ER Visits Age-Adjusted Rate per 10,000 population, 2015	Percentage of live births to mothers who smoked during pregnancy, 2017	Estimated cost of smoking-related births, 2017
Adams	19%	42.7	219.3	21.9	9.2	\$86,912
Allen	21%	43.5	225.1	45.0	10.3	\$726,530
Bartholomew	24%	47.5	242.7	40.5	12.6	\$179,256
Benton	Suppressed	50.5	260.1	26.8	20.2	\$29,876
Blackford	24%	69.7	244.6	43.3	40.0	\$67,900
Boone	15%	48.6	257.3	24.4	7.4	\$77,406
Brown	18%	41.4	196	Unstable Rate	17.5	\$28,518
Carroll	23%	47.7	207.1	24.2	14.3	\$46,172
Cass	33%	54.6	232	43.6	15.1	\$93,702
Clark	23%	62.8	270.7	25.6	11.6	\$232,218
Clay	18%	67.7	299.8	35.9	16.1	\$69,258
Clinton	23%	50.1	259.4	40.8	17.7	\$103,208
Crawford	37%	73.6	251.8	24.6	30.6	\$50,246
Daviess	13%	47.7	257.1	47.2	12.7	\$95,060
Dearborn	23%	57.5	226.8	25.5	17.8	\$115,430
Decatur	16%	48.3	255.9	49.2	17.1	\$74,690
DeKalb	28%	50.7	260.7	26.0	18.8	\$130,368
Delaware	21%	53.3	257.1	45.0	22.6	\$344,932
Dubois	13%	31.9	246.1	5.6	8.4	\$65,184
Elkhart	21%	41.7	235.8	44.5	9.3	\$392,462
Fayette	31%	57.6	282.2	27.5	21.2	\$80,122
Floyd	18%	52.3	263.6	27.1	10.3	\$123,578
Fountain	25%	48.5	232.8	60.9	18.4	\$46,172
Franklin	18%	45.5	224	12.4	15.7	\$59,752
Fulton	13%	59.4	268.7	35.0	25.6	\$78,764
Gibson	16%	47.8	243.8	47.4	20.3	\$108,640
Grant	32%	59.9	269.6	59.1	31.2	\$329,994
Greene	30%	63.6	270.1	24.8	20.3	\$96,418
Hamilton	10%	29.7	174.1	19.5	2.3	\$122,220
Hancock	16%	52.7	205.4	29.0	9.1	\$104,566
Harrison	19%	61.4	241.1	23.0	16.6	\$96,418
Hendricks	14%	45.6	203.6	15.3	7.4	\$173,824
Henry	23%	54.7	250.4	46.1	24.7	\$150,738
Howard	28%	52.3	273.3	57.3	20.8	\$274,316
Huntington	27%	43.3	271.7	40.0	21.5	\$123,578
Jackson	21%	68.3	230.7	67.9	23.1	\$186,046
Jasper	25%	48.8	267.1	34.1	16.0	\$81,480
Jay	22%	61.3	260.8	54.2	16.0	\$62,468
Jefferson	34%	72.2	312	31.0	28.2	\$143,948
Jennings	33%	68.8	283.8	55.6	28.0	\$131,726
Johnson	21%	47.6	226.6	39.0	12.5	\$309,624
Knox	18%	54.5	273.1	43.6	22.4	\$126,294
Kosciusko	25%	47.7	232.8	28.7	15.1	\$217,280
LaGrange	20%	38.7	239.1	27.4	6.2	\$63,826
Lake	22%	47.9	253.1	69.9	8.4	\$643,692
LaPorte	29%	52	279.4	52.5	21.6	\$373,450

(Continued on next page)

APPENDIX 3D - Part 1 (Continued from previous page)

County	Estimated adult smoking rate (Statewide: 2017 BRFSS; County-level: 2013-2017 BRFSS)	Age-adjusted rate of lung cancer deaths per 100,000 population (2013-2017)	Age-adjusted rate of major cardiovascular diseases deaths per 100,000 population (2013-2017)	Asthma ER Visits Age-Adjusted Rate per 10,000 population, 2015	Percentage of live births to mothers who smoked during pregnancy, 2017	Estimated cost of smoking-related births, 2017
Lawrence	28%	61.6	269	50.5	28.1	\$183,330
Madison	28%	60.6	245.4	87.0	20.1	\$382,956
Marion	23%	55.7	244.7	83.4	10.5	\$2,039,716
Marshall	27%	46.9	224.6	25.9	16.3	\$134,442
Martin	18%	57.3	244.8	Unstable Rate	17.7	\$27,160
Miami	34%	49.1	324	45.0	24.1	\$129,010
Monroe	20%	41	187	22.9	15.4	\$262,094
Montgomery	21%	50.2	267.9	51.3	19.0	\$116,788
Morgan	21%	59.4	255.1	41.6	23.7	\$259,378
Newton	44%	67.9	231.1	31.1	22.3	\$42,098
Noble	25%	50.2	226.2	32.8	16.7	\$142,590
Ohio	Suppressed	62.8	202.5	Unstable Rate	12.9	\$12,222
Orange	23%	56.3	264.8	52.8	23.6	\$81,480
Owen	32%	68	276.2	32.8	23.3	\$65,184
Parke	27%	54.9	257.7	32.3	15.1	\$42,098
Perry	24%	49.7	268.1	73.8	28.5	\$82,838
Pike	Suppressed	56.6	250.1	Unstable Rate	16.7	\$29,876
Porter	18%	45.5	209.3	44.0	10.7	\$239,008
Posey	24%	53.1	214.3	20.7	22.7	\$78,764
Pulaski	24%	53.6	292.5	29.4	27.1	\$47,530
Putnam	28%	65	241.4	25.1	19.3	\$96,418
Randolph	18%	50.7	238.2	47.7	22.6	\$89,628
Ripley	24%	52.8	255.9	39.0	19.2	\$88,270
Rush	19%	63	255.6	83.1	22.8	\$58,394
Scott	31%	75	288	51.9	25.1	\$93,702
Shelby	21%	62.1	236.4	51.6	21.2	\$143,948
Spencer	12%	48.4	233.3	22.5	12.1	\$35,308
St. Joseph	22%	47.3	239.1	50.6	10.6	\$505,176
Starke	27%	76.2	314.2	51.8	21.6	\$77,406
Steuben	23%	50.8	219.4	40.7	21.0	\$97,776
Sullivan	18%	67.8	282.5	46.8	20.4	\$59,752
Switzerland	Suppressed	47.9	251.9	Unstable Rate	20.4	\$31,234
Tippecanoe	21%	44.3	228.2	38.0	10.5	\$319,130
Tipton	17%	48.4	222.7	40.4	16.3	\$32,592
Union	Suppressed	Unreliable	257.2	Suppressed	14.5	\$13,580
Vanderburgh	21%	51	224.6	54.9	17.4	\$503,818
Vermillion	30%	53.8	366.9	48.7	22.4	\$51,604
Vigo	24%	60.6	287.6	44.9	21.3	\$354,438
Wabash	20%	43	241.2	27.4	26.3	\$112,714
Warren	43%	42.4	219.8	47.3	14.0	\$16,296
Warrick	15%	42.6	201.8	30.1	10.9	\$95,060
Washington	26%	66.6	289.5	44.3	20.5	\$89,628
Wayne	22%	57.5	290.5	41.9	14.5	\$153,454
Wells	17%	44.8	220.8	28.0	18.6	\$82,838
White	15%	49.6	235.5	53.8	18.7	\$76,048
Whitley	18%	48.3	221.7	35.1	14.1	\$78,764
Indiana	21.8%	50.7	243.9	47.4	13.5	\$15,073,800

Source: ISDH/TPCC, 2019

APPENDIX 3D - Part 2

County	Estimated number of people living with a tobacco-related illness	Estimated number of deaths due to tobacco	Estimated number of deaths due to secondhand smoke (SHS)	Estimated cost of SHS due to medical costs and premature death
Adams	1,617	54	7	\$11.5 Million
Allen	17,715	591	73	\$118.7 Million
Bartholomew	3,923	131	16	\$25.7 Million
Benton	449	15	2	\$3 Million
Blackford	673	22	3	\$4.3 Million
Boone	2,781	93	12	\$18.9 Million
Brown	824	27	3	\$5.1 Million
Carroll	1,038	35	4	\$6.7 Million
Cass	1,972	66	8	\$13 Million
Clark	5,746	192	23	\$36.8 Million
Clay	1,397	47	6	\$9 Million
Clinton	1,665	55	7	\$11.1 Million
Crawford	561	19	2	\$3.6 Million
Daviess	1,539	51	7	\$10.6 Million
Dearborn	2,563	85	10	\$16.7 Million
Decatur	1,310	44	5	\$8.6 Million
DeKalb	2,123	71	9	\$14.1 Million
Delaware	6,427	214	24	\$39.3 Million
Dubois	2,132	71	9	\$14 Million
Elkhart	9,657	322	41	\$66 Million
Fayette	1,261	42	5	\$8.1 Million
Floyd	3,869	129	15	\$24.9 Million
Fountain	892	30	4	\$5.8 Million
Franklin	1,165	39	5	\$7.7 Million
Fulton	1,070	36	4	\$7 Million
Gibson	1,732	58	7	\$11.2 Million
Grant	3,749	125	14	\$23.4 Million
Greene	1,727	58	7	\$11.1 Million
Hamilton	13,089	436	57	\$91.7 Million
Hancock	3,529	118	14	\$23.4 Million
Harrison	2,053	68	8	\$13.1 Million
Hendricks	7,208	240	30	\$48.6 Million
Henry	2,624	87	10	\$16.5 Million
Howard	4,314	144	17	\$27.6 Million
Huntington	1,935	64	8	\$12.4 Million
Jackson	2,183	73	9	\$14.2 Million
Jasper	1,700	57	7	\$11.2 Million
Jay	1,066	36	4	\$7.1 Million
Jefferson	1,714	57	7	\$10.8 Million
Jennings	1,434	48	6	\$9.5 Million
Johnson	7,018	234	29	\$46.6 Million
Knox	2,066	69	8	\$12.8 Million
Kosciusko	3,930	131	16	\$25.8 Million
LaGrange	1,661	55	8	\$12.4 Million
Lake	25,185	839	102	\$165.7 Million
LaPorte	5,880	196	23	\$37.2 Million
Lawrence	2,408	80	10	\$15.4 Million
Madison	6,915	231	27	\$44 Million
Marion	46,232	1,541	186	\$301.8 Million
Marshall	2,350	78	10	\$15.7 Million
Martin	536	18	2	\$3.5 Million

(Continued on next page)

APPENDIX 3D - Part 2

(Continued from previous page)

County	Estimated number of people living with a tobacco-related illness	Estimated number of deaths due to tobacco	Estimated number of deaths due to secondhand smoke (SHS)	Estimated cost of SHS due to medical costs and premature death
Miami	1,947	65	8	\$12.3 Million
Monroe	7,889	263	28	\$46.1 Million
Montgomery	1,980	66	8	\$12.7 Million
Morgan	3,522	117	14	\$23 Million
Newton	749	25	3	\$4.8 Million
Noble	2,369	79	10	\$15.9 Million
Ohio	330	11	1	\$2 Million
Orange	1,021	34	4	\$6.6 Million
Owen	1,131	38	4	\$7.2 Million
Parke	931	31	4	\$5.8 Million
Perry	1,038	35	4	\$6.5 Million
Pike	681	23	3	\$4.3 Million
Porter	8,498	283	34	\$54.9 Million
Posey	1,350	45	5	\$8.7 Million
Pulaski	697	23	3	\$4.5 Million
Putnam	2,047	68	8	\$12.7 Million
Randolph	1,352	45	5	\$8.7 Million
Ripley	1,450	48	6	\$9.6 Million
Rush	894	30	4	\$5.8 Million
Scott	1,255	42	5	\$8.1 Million
Shelby	2,294	76	9	\$14.8 Million
Spencer	1,085	36	4	\$7 Million
St. Joseph	13,734	458	55	\$89.2 Million
Starke	1,207	40	5	\$7.8 Million
Steuben	1,800	60	7	\$11.4 Million
Sullivan	1,153	38	4	\$7.2 Million
Switzerland	539	18	2	\$3.5 Million
Tippecanoe	9,361	312	36	\$57.7 Million
Tipton	836	28	3	\$5.3 Million
Union	385	13	2	\$2.5 Million
Vanderburgh	9,549	318	37	\$60 Million
Vermillion	852	28	3	\$5.4 Million
Vigo	5,792	193	22	\$36 Million
Wabash	1,737	58	7	\$11 Million
Warren	445	15	2	\$2.8 Million
Warrick	3,023	101	12	\$19.9 Million
Washington	1,444	48	6	\$9.4 Million
Wayne	3,622	121	14	\$23 Million
Wells	1,416	47	6	\$9.2 Million
White	1,276	43	5	\$8.2 Million
Whitley	1,715	57	7	\$11.1 Million
Indiana	333,000	11,100	1,337	\$2.1 Billion

Source: ISDH/TPCC, 2019

REFERENCES, CHAPTER 3

- Campaign for Tobacco-Free Kids. (2018a). *State cigarette excise tax rates & rankings*. Retrieved from <http://www.tobaccofreekids.org/research/factsheets/pdf/0097.pdf>
- Campaign for Tobacco-Free Kids. (2018b). *The toll of tobacco in Indiana*. Retrieved from http://www.tobaccofreekids.org/facts_issues/toll_us/indiana
- Centers for Disease Control and Prevention. (1991-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (2016a). *Health effects of cigarette smoking*. Retrieved from https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/
- Centers for Disease Control and Prevention. (2016b). *Youth and tobacco use*. Retrieved from https://www.cdc.gov/tobacco/data_statistics/fact_sheets/youth_data/tobacco_use/index.htm
- Centers for Disease Control and Prevention. (2017). *Behavioral Risk Factor Surveillance System (BRFSS) prevalence & trends data*. Retrieved from <http://www.cdc.gov/brfss/brfssprevalence/index.html>
- Centers for Disease Control and Prevention. (2018a). *Economic Trends in Tobacco*. Retrieved from https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/
- Centers for Disease Control and Prevention. (2018b). *Quitting Smoking*. Retrieved from https://www.cdc.gov/tobacco/data_statistics/fact_sheets/cessation/quitting/index.htm
- Federal Trade Commission. (2018). *Federal Trade Commission cigarette report for 2016 and Federal Trade Commission smokeless tobacco report for 2016*. Retrieved from https://www.ftc.gov/system/files/documents/reports/federal-trade-commission-cigarette-report-2016-federal-trade-commission-smokeless-tobacco-report/ftc_cigarette_report_for_2016_0.pdf
- Gassman, R., Jun, M., Samuel, S., Agle, J. D., King, R., Ables, E.,... Wolf, J. (2018). *Indiana Youth Survey*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Indiana Department of Education. (2019). *DOE discipline 2018, incidents and student counts*. Aggregate data received from Jeff Wittman on January 18, 2019.
- Indiana State Department of Health, Epidemiology Resource Center. (2018). *Indiana natality report 2017 (Table 20)*. Retrieved from <http://www.in.gov/isdh/19095.htm>
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2017). *Health effects of secondhand smoke*. Retrieved from https://www.in.gov/isdh/tpc/files/Health%20Effects%20of%20Secondhand%20Smoke_10_06_2017.pdf
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2018a). *Electronic cigarettes*. Retrieved from https://www.in.gov/isdh/tpc/files/Electronic%20Cigarettes_%2010_19_2018.pdf
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2018b). *Indiana adult smoking*. Retrieved from https://www.in.gov/isdh/tpc/files/IN%20Adult%20Smoking_10_19_2018.pdf
- Indiana State Department of Health, Tobacco Prevention and Cessation Commission. (2019). *Indiana Youth Tobacco Survey; Indiana Adult Tobacco Survey; Indiana County-level Tobacco Use & Outcomes*. Data received on January 25, 2019, from Katelin Rupp, Director of Program Evaluation at the Indiana State Department of Health, Tobacco Prevention & Cessation Commission.
- Inter-university Consortium for Political and Social Research, University of Michigan. (2018). *Monitoring the Future (MTF)*. Retrieved from <http://www.monitoringthefuture.org/data/data.html>
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey*. Bloomington, IN: Institute for Research on Addictive Behavior, Indiana University. Retrieved from <https://iprc.iu.edu/indiana-college-survey/substance-use-survey>

- Lewis, C., & Zollinger, T. (2014). *Estimating the Economic Impact of Secondhand Smoke in Indiana in 2014*. Retrieved from [http://www.in.gov/isdh/tpc/files/Executive%20Summary%20Estimating%20the%20Economic%20Impact%20of%20%20Secondhand%20Smoke%20in%20%20%20%20\(002\).pdf](http://www.in.gov/isdh/tpc/files/Executive%20Summary%20Estimating%20the%20Economic%20Impact%20of%20%20Secondhand%20Smoke%20in%20%20%20%20(002).pdf)
- Marynak, K., Gammon, D., King, B., Loomis, B., Fulmer, E., Wang, T., & Rogers, T. (2017). National and State Trends in Sales of Cigarettes and E-Cigarettes, U.S., 2011–2015. *American Journal of Preventive Medicine* (in press). Retrieved from <http://www.sciencedirect.com/science/article/pii/S0749379717300223>
- Siegel, R. L., Miller, K. D., & Jemal, A. (2015). Cancer Statistics, 2015. *CA: A cancer journal for clinicians*, 65(1), 5-29.
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>
- U.S. Department of Health and Human Services. (2014). *The Health Consequences of Smoking -- 50 years of progress: A Report of the Surgeon General*. Atlanta, GA.: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health.
- U.S. Department of Health and Human Services. (2016). *E-cigarette use among youth and young adults: A Report of the Surgeon General*. Atlanta, GA.: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health.
- World Health Organization. (2015). *WHO global report on trends in tobacco smoking 2000-2025*. Retrieved from <http://www.who.int/tobacco/publications/surveillance/reportontrendstobaccosmoking/en/>

4 MARIJUANA USE IN INDIANA: CONSUMPTION PATTERNS AND CONSEQUENCES

INTRODUCTION

Marijuana is a product of the hemp plant, known as *Cannabis sativa*. Found in the dried leaves, stems, seeds, and flowers, delta-9-tetrahydrocannabinol (THC) is the primary psychoactive (mind-altering) chemical. The drug can be consumed in different ways: by smoking “joints” or “blunts” (hand-rolled cigarettes or cigars filled only with cannabis, not tobacco) and hookahs (water pipes), mixing into foods, or brewing as tea (Hall & Solowij, 1998). Recent studies show an increase in edible consumption of marijuana, especially in states that allow medical use of marijuana (National Institute on Drug Abuse [NIDA], 2016a). Marijuana is the most commonly used illicit drug in the United States (Azofeifa et al., 2016).

Age of first use is an important risk factor in the subsequent progression to substance misuse and dependence (King & Chassin, 2007). Adolescents who used marijuana by the age of 17 were found to be at greater risk of using other drugs and developing alcohol and drug abuse/dependence (Lynskey et al., 2003). The use of marijuana can result in adverse physical, mental, emotional, and behavioral changes. Short-term effects include memory impairment and learning problems, distorted perception, difficulty thinking and solving problems, loss of coordination, and increased heart rate. Long-term use has been linked to respiratory illnesses

and an increased risk of heart attack and cancer (Crean, Crane, & Mason, 2011; Volkow, Baler, Compton, & Weiss, 2014). Furthermore, prolonged marijuana use can lead to mental health problems such as depression, anxiety, suicidal thoughts, and personality disturbances (Patton et al., 2002; Caspi et al., 2005).


Babies born to women who used marijuana during their pregnancy may be at an increased risk for neurobehavioral problems, potentially exhibiting difficulties with attention, memory, and problem solving (NIDA, 2016a).

PREVALENCE OF MARIJUANA CONSUMPTION IN THE GENERAL POPULATION

National Survey on Drug Use and Health

According to the 2017 National Survey on Drug Use and Health (NSDUH), an estimated 9.3% (95% Confidence Interval [CI]: 8.0–10.8) of Indiana residents ages 12 and older reported current (past-month) marijuana use (U.S.: 9.2%; 95% CI: 9.0–9.5); past-year use was estimated at 14.4% (95% CI: 12.8–16.2; U.S.: 14.5%; 95% CI: 14.2–14.8) (Substance Abuse and Mental Health Services Administration [SAMHSA], 2017). For 10-year trend data on past-month marijuana use, see Figure 4.1.

Figure 4.1 Percentage of Indiana and U.S. Population (Ages 12 and Older) Reporting Current Marijuana Use (National Survey on Drug Use and Health, 2008–2017)




Source: SAMHSA, 2017

The highest prevalence was among individuals ages 18 to 25, with 20.7% (95% CI: 17.6–24.2) of Hoosiers in this age group reporting current marijuana use (U.S.: 21.5%; 95% CI: 20.8–22.1) and 33.9% (95% CI: 30.1–37.8) reporting past-year use (U.S.: 33.9%; 95% CI: 33.3–34.7) in 2017 (Figure 4.2). Prevalence rates were significantly lower in youth and adults ages 26 and older. Based on 2017 estimates, 13.3% (95% CI: 11.1–15.9) of 12- to 17-year-olds in Indiana reported using marijuana

in the past year (U.S.: 12.2%; 95% CI: 11.8–12.6) and 6.5% (95% CI: 5.2–8.1) in the past month (U.S.: 6.5%; 95% CI: 6.1–6.8). Among Hoosiers ages 26 and older, 7.7% (95% CI: 6.2–9.4) reported past-month marijuana use (U.S.: 7.6%; 95% CI: 7.3–7.8) and 11.2% (95% CI: 9.5–13.2) reported use in the past year (U.S.: 11.6%; 95% CI: 11.3–12.0) (SAMSHA, 2017). See Figure 4.2 for current marijuana use rates by age group in Indiana.

Figure 4.2 Percentage of Indiana Residents Reporting Current Marijuana Use, by Age Group (National Survey on Drug Use and Health, 2008–2017)




Source: SAMHSA, 2017

Marijuana initiation, or first-time use, was primarily reported in young adults and adolescents. An estimated 8.8% (95% CI: 7.1–11.0) of Hoosiers ages 18 to 25 initiated marijuana use in the past year (U.S.: 8.0%; 95% CI: 7.6–8.4), as did 6.0% (95% CI: 5.0–7.3) of Indiana youth ages 12 to 17 (U.S.: 5.2%; 95% CI: 5.0–5.5). Past-year initiation was significantly lower in adults ages 26 and older (IN: 0.5%; 95% CI: 0.4–0.7; U.S.: 0.5%; 95% CI: 0.4–0.5) (SAMHSA, 2017).

Youth Risk Behavior Surveillance System

The Youth Risk Behavior Surveillance System (YRBSS) estimated that in 2015, 16.4% (95% CI: 14.1–18.9) of Indiana high school students used marijuana in the past month; this percentage is significantly lower than the national rate of 21.7% (95% CI: 19.3–24.2). Use was more likely to occur in higher grade levels and in black or Hispanic students (Centers for Disease Control and Prevention [CDC], 1999–2015). For more detailed information, see Table 4.1 and Figure 4.3.

Figure 4.3 Percentage of Indiana and U.S. High School Students Currently Using Marijuana (Youth Risk Behavior Surveillance System, 2003–2015)



Note: 2013 estimates are not available for Indiana.
Source: CDC, 1999–2015

Table 4.1 Percentage of Indiana and U.S. High School Students Reporting Current (Past Month) Marijuana Use, by Grade, Gender, and Race/Ethnicity (Youth Risk Behavior Surveillance System, 2015)

		Indiana (95% CI)	U.S. (95% CI)
Grade	9th	13.7% (10.4–17.9)	15.2% (16.8–23.5)
	10th	16.8% (12.5–22.2)	20.0% (24.0–30.4)
	11th	17.0% (13.2–21.7)	24.8% (22.3–27.5)
	12th	18.4% (14.1–23.7)	27.6% (23.8–31.6)
Gender	Male	16.4% (13.8–19.4)	23.2% (20.4–26.3)
	Female	15.9% (12.7–19.7)	20.1% (17.6–22.9)
Race/Ethnicity	Black	23.2% (17.1–30.7)	28.9% (26.3–31.6)
	White	14.9% (12.4–17.8)	20.4% (17.8–23.3)
	Hispanic	18.1% (13.6–23.6)	27.6% (24.6–30.7)
Total		16.4% (14.1–18.9)	21.7% (19.3–24.2)


Source: CDC, 2017

In 2015, 6.2% (95% CI: 5.3–7.4) of Indiana students reported having tried marijuana before the age of 13; that figure was comparable to the national rate (7.5%; 95% CI: 6.5–8.7) (CDC, 1999–2015).

Indiana Youth Survey

Data from the Indiana Youth Survey (Gassman et al., 2018), and the Monitoring the Future (MTF) survey (Inter-university Consortium for Political and Social Research [ICPSR], 2018) show that marijuana use among 8th, 10th, and 12th grade students increased with grade level/age. Prevalence rates for current marijuana use in Indiana and the nation were similar; however, due to lack of detail in the publicly available dataset, statistical significance could not be determined. For current marijuana use trends among 8th, 10th, and 12th grade students from 2009 through 2018, see Figure 4.4; for monthly marijuana use by Indiana region and grade level for 2018, see Appendix 4A.

Figure 4.4 Percentage of Indiana and U.S. 8th, 10th, and 12th Grade Students Reporting Current Marijuana Use (Indiana Youth Survey and Monitoring the Future Survey, 2009–2018)



Source: Gassman et al., 2018; ICPSR, 2018

Indiana College Substance Use Survey


Marijuana use was also prevalent among college students. Results from the 2018 Indiana College Substance Use Survey showed that 22.7% of Indiana college students reported current marijuana use (U.S.: 21.2%). Users were more likely to be male and between the ages of 21 and 25 (King & Jun, 2018).¹

USE OF MARIJUANA IN THE TREATMENT POPULATION

Treatment Episode Data Set

The Treatment Episode Data Set (TEDS) collects information from clients being admitted to substance abuse treatment. The data show that from 2007 through 2016, Indiana exhibited a significantly higher percentage of treatment episodes reporting marijuana use and dependence² compared to the rest of the United States. From 2007 through 2016, roughly one-half of Indiana treatment admissions reported marijuana use and about one-fifth indicated marijuana dependence (see Figure 4.5) (SAMHSA, 2016).

Figure 4.5 Percentage of Indiana and U.S. Treatment Episodes with Marijuana Use and Marijuana Dependence Reported at Treatment Admission (Treatment Episode Data Set, 2007–2016)



Source: SAMHSA, 2016

¹Twenty-four Indiana colleges participated in the survey; results are based on nonrandom sampling and are not representative of all college students in Indiana.

²We defined marijuana dependence as “individuals in substance abuse treatment listing marijuana as their primary substance at admission.”

Table 4.2 Percentage of Indiana Treatment Admissions with Reported Marijuana Use, by Gender, Race, and Age Group (Treatment Episode Data Set, 2016)

		Marijuana Use
Gender	Male	51.5%
	Female	43.7%
Race	White	46.6%
	Black	57.4%
	Other	52.7%
Age Group	Under 18	89.9%
	18-24	66.4%
	25-34	48.9%
	35-44	41.7%
	45-54	32.9%
	55+	24.8%
Total		48.4%

Source: SAMHSA, 2016

Statistically significant differences in marijuana use among Indiana’s treatment population were observed by gender, race, and age, as follows:

- The percentage of males reporting marijuana use was higher than the percentage of females.
- The percentage of Blacks who reported marijuana use was higher compared to Whites and other races.
- Marijuana use decreased by age; i.e., the highest percentage was found among adolescents under the age of 18 and the lowest among older adults ages 55 and above (see Table 4.2).

For county-level information on marijuana use and dependence, see Appendix 4B (Indiana Family and Social Services Administration, 2018).

CONSEQUENCES OF MARIJUANA USE

The debate around the potential harms and benefits of marijuana use is perhaps the most fervent it has ever been as numerous states have or are in the process of implementing medical and recreational marijuana legislation. As of December 2018, 33 states and the District of Columbia (D.C.) have legalized medical marijuana use, and 10 of these states, as well as D.C., have passed laws to allow adult recreational use (National Conference of State Legislatures, 2018). Existing research shows that marijuana use is associated with numerous deleterious health effects. Short-term use is associated with impaired motor coordination and altered judgement that can increase the likelihood of other risky behaviors. Long-term use can increase the risk of mental illness, use of other substances, and chronic bronchitis (Volkow et al., 2014).

Persistent cannabis use is associated with decreased functional connectivity in the brain, IQ decline, and increased memory and attention issues (Meier et al., 2012; Zalesky et al., 2012). In addition to the aforementioned negative health effects, cannabis dependence can have undesirable economic and social implications. A longitudinal study found that regular users of cannabis were more likely to end up in a lower social class than their parents, have more relationship and workplace problems, and experience more financial difficulties in early midlife (Cerdá et al., 2016). Conversely, medical marijuana use has been shown to relieve the symptoms of numerous clinical conditions including glaucoma, nausea, chronic pain, inflammation, disease-induced decreased appetite, multiple sclerosis, and epilepsy (Volkow et al., 2014).

APPENDIX 4A

Percentage of Indiana Students Reporting Monthly Marijuana Use, by Region and Grade (Indiana Youth Survey, 2018)

	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
6th Grade	0.9	0.7	0.8	1.4	1.3	1.2	1.1	0.6	0.7
7th Grade	2.9	2.9	2.7	1.9	1.9*	4.7*	3.1	1.6*	3.0
8th Grade	5.9	6.9*	5.8	8.2*	4.6*	7.8*	7.2*	3.9*	5.2
9th Grade	8.9	11.9*	9.1	7.0*	8.1	7.8	9.9	7.8	7.6*
10th Grade	12.6	14.6*	11.7	9.1*	12.7	13.7	14.4*	11.6	11.8
11th Grade	15.1	18.9*	13.4	11.0*	16.4	16.0	16.0	14.7	13.1*
12th Grade	17.3	20.5*	15.1*	11.2	18.0	16.3	18.9	13.9*	18.7

Note: * Indicates a regional rate that is significantly different from the overall state rate ($P < 0.05$).

Source: Gassman et al., 2018

APPENDIX 4B

Number of Treatment Admissions with Marijuana Use and Dependence Reported at Treatment Admission in Indiana, by County (Substance Abuse Population by County/Treatment Episode Data Set, 2018)

County	Treatment Episodes	Marijuana Use		Marijuana Dependence	
	Total	Number	%	Number	%
Adams	228	139	61.0%	66	28.9%
Allen	2,050	1,158	56.5%	545	26.6%
Bartholomew	727	334	45.9%	91	12.5%
Benton	61	43	70.5%	16	26.2%
Blackford	144	83	57.6%	18	12.5%
Boone	167	83	49.7%	32	19.2%
Brown	72	33	45.8%	8	11.1%
Carroll	80	35	43.8%	12	15.0%
Cass	387	202	52.2%	88	22.7%
Clark	103	31	30.1%	11	10.7%
Clay	106	37	34.9%	17	16.0%
Clinton	147	64	43.5%	26	17.7%
Crawford	36	16	44.4%	5	13.9%
Daviess	202	78	38.6%	30	14.9%
Dearborn	497	246	49.5%	77	15.5%
Decatur	185	96	51.9%	42	22.7%
DeKalb	276	152	55.1%	50	18.1%
Delaware	843	314	37.2%	92	10.9%
Dubois	131	68	51.9%	31	23.7%
Elkhart	765	404	52.8%	178	23.3%
Fayette	322	131	40.7%	54	16.8%
Floyd	49	10	20.4%	5	10.2%
Fountain	59	35	59.3%	14	23.7%
Franklin	145	70	48.3%	28	19.3%
Fulton	188	99	52.7%	23	12.2%
Gibson	252	135	53.6%	59	23.4%
Grant	509	247	48.5%	83	16.3%
Greene	197	107	54.3%	38	19.3%
Hamilton	874	519	59.4%	268	30.7%
Hancock	398	200	50.3%	109	27.4%
Harrison	19	<5	N/A	<5	N/A
Hendricks	424	187	44.1%	96	22.6%
Henry	381	151	39.6%	73	19.2%
Howard	492	206	41.9%	58	11.8%
Huntington	136	73	53.7%	32	23.5%
Jackson	330	196	59.4%	40	12.1%
Jasper	93	37	39.8%	17	18.3%
Jay	122	49	40.2%	15	12.3%
Jefferson	307	162	52.8%	76	24.8%
Jennings	260	127	48.8%	71	27.3%
Johnson	424	161	38.0%	54	12.7%
Knox	352	188	53.4%	74	21.0%
Kosciusko	362	206	56.9%	76	21.0%
LaGrange	115	65	56.5%	29	25.2%
Lake	1,903	743	39.0%	317	16.7%
LaPorte	172	61	35.5%	16	9.3%
Lawrence	535	295	55.1%	91	17.0%

County	Treatment Episodes	Marijuana Use		Marijuana Dependence	
	Total	Number	%	Number	%
Madison	1,205	676	56.1%	336	27.9%
Marion	4,134	1,856	44.9%	939	22.7%
Marshall	137	77	56.2%	40	29.2%
Martin	60	29	48.3%	10	16.7%
Miami	249	126	50.6%	42	16.9%
Monroe	1,428	788	55.2%	239	16.7%
Montgomery	304	163	53.6%	84	27.6%
Morgan	550	253	46.0%	100	18.2%
Newton	27	10	37.0%	<5	N/A
Noble	346	186	53.8%	86	24.9%
Ohio	33	14	42.4%	6	18.2%
Orange	135	62	45.9%	27	20.0%
Owen	187	91	48.7%	30	16.0%
Parke	65	36	55.4%	19	29.2%
Perry	94	40	42.6%	12	12.8%
Pike	46	26	56.5%	11	23.9%
Porter	202	81	40.1%	32	15.8%
Posey	156	86	55.1%	42	26.9%
Pulaski	82	38	46.3%	9	11.0%
Putnam	253	127	50.2%	51	20.2%
Randolph	167	72	43.1%	24	14.4%
Ripley	180	70	38.9%	28	15.6%
Rush	139	72	51.8%	27	19.4%
Saint Joseph	1,351	606	44.9%	281	20.8%
Scott	212	85	40.1%	20	9.4%
Shelby	140	69	49.3%	14	10.0%
Spencer	132	71	53.8%	21	15.9%
Starke	77	27	35.1%	5	6.5%
Steuben	183	88	48.1%	31	16.9%
Sullivan	115	50	43.5%	21	18.3%
Switzerland	113	58	51.3%	21	18.6%
Tippecanoe	364	199	54.7%	94	25.8%
Tipton	46	29	63.0%	21	45.7%
Union	54	13	24.1%	<5	N/A
Vanderburgh	1,218	724	59.4%	325	26.7%
Vermillion	132	65	49.2%	26	19.7%
Vigo	771	432	56.0%	227	29.4%
Wabash	299	169	56.5%	62	20.7%
Warren	26	13	50.0%	9	34.6%
Warrick	310	193	62.3%	92	29.7%
Washington	44	19	43.2%	<5	N/A
Wayne	558	263	47.1%	91	16.3%
Wells	156	109	69.9%	53	34.0%
White	125	66	52.8%	17	13.6%
Whitley	126	71	56.3%	26	20.6%
Indiana	33,399	16,489	49.4%	6,910	20.7%

Note: We defined marijuana dependence as “individuals in substance abuse treatment listing marijuana as their primary substance at admission.”

We calculated the percentages by dividing the number of reported marijuana use/dependence by the number of treatment episodes.

Information on treatment episodes <5 was suppressed due to confidentiality constraints.

Source: Indiana Family and Social Services Administration, 2018

REFERENCES, CHAPTER 4

- Azofeifa, A., Mattson, M. E., Schauer, G., McAfee, T., Grant, A., & Lyerla, R. (2016). National Estimates of Marijuana Use and Related Indicators—National Survey on Drug Use and Health, United States, 2002–2014. *MMWR. Surveillance Summaries*, *65*, 1-25.
- Caspi, A., Moffitt, T. E., Cannon, M., McClay, J., Murray, R., Harrington, H., ... Craig, I. W. (2005). Moderation of the effect of adolescent-onset cannabis use on adult psychosis by a functional polymorphism in the catechol-O-methyltransferase gene: Longitudinal evidence of a gene X environment interaction. *Biological Psychiatry*, *57*, 1117-1127.
- Centers for Disease Control and Prevention. (1999-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Cerdá, M., Moffitt, T. E., Meier, M. H., Harrington, H., Houts, R., Ramrakha, S., ... & Caspi, A. (2016). Persistent cannabis dependence and alcohol dependence represent risks for midlife economic and social problems: a longitudinal cohort study. *Clinical psychological science*, *4*(6), 1028-1046.
- Crean, R. D., Crane, N.A., & Mason, B. J. (2011). An evidence based review of acute and long-term effects of cannabis use on executive cognitive functions. *Journal of Addiction Medicine*, *5*, 1-8.
- Federal Bureau of Investigation. (2014). *Uniform Crime Reporting (UCR) program data: County-level detailed arrest and offense data, 2012*. ICPSR35019-v1. Ann Arbor, MI: Inter-university Consortium for Political and Social Research [distributor], 2014-06-12. Retrieved from <http://doi.org/10.3886/ICPSR35019.v1>
- Gassman, R., Jun, M., Samuel, S., Agle, J. D., King, R., Ables, E., ... Wolf, J. (2018). *Indiana Youth Survey-2018*. Institute for Research on Addictive Behavior, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Hall, W., & Solowij, N. (1998). Adverse effects of cannabis. *The Lancet*, *352*, 1611-1616.
- Indiana Family and Social Services Administration. (2018). *Treatment Episode Data System (TEDS)*. Indianapolis, IN: Indiana Family and Social Services Administration.
- Inter-university Consortium for Political and Social Research, University of Michigan. (2018). *Monitoring the Future (MTF)*. Retrieved from <http://www.monitoringthefuture.org/data/data.html>
- King, K. M., & Chassin, L. (2007). A prospective study of the effects of age of initiation of alcohol and drug use on young adult substance dependence. *Journal of Studies on Alcohol and Drugs*, *68*, 256-265.
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey, 2018*. Indiana University. Retrieved from <http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey>
- Lynskey, M. T., Heath, A. C., Bucholz, K. K., Slutske, W. S., Madden, P. A., Nelson, E. C., ... Martin, N. G. (2003). Escalation of drug use in early-onset cannabis users vs. co-twin controls. *Journal of the American Medical Association*, *289*, 427-433.
- Meier, M. H., Caspi, A., Ambler, A., Harrington, H., Houts, R., Keefe, R. S., ... & Moffitt, T. E. (2012). Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proceedings of the National Academy of Sciences*, 201206820.
- National Conference of State Legislatures. (2018). *Marijuana overview*. Retrieved from <http://www.ncsl.org/research/civil-and-criminal-justice/marijuana-overview.aspx>
- National Institute on Drug Abuse. (2016a). *DrugFacts: Marijuana*. National Institutes of Health, United States Department of Health and Human Services. Retrieved from <http://www.drugabuse.gov/publications/drugfacts/marijuana>

Patton, G. C., Coffey, C., Carlin, J. B., Degenhardt, L., Lynskey, M., & Hall, W. (2002). Cannabis use and mental health in young people: cohort study. *British Medical Journal*, *325*, 1195-1198.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>

Volkow, N. D., Baler, R. D., Compton, W. M., & Weiss, S. R. B. (2014). Adverse health effects of marijuana use. *The New England Journal of Medicine*, *370*, 2219-2227.

Zalesky, A., Solowij, N., Yücel, M., Lubman, D. I., Takagi, M., Harding, I. H., ... & Seal, M. (2012). Effect of long-term cannabis use on axonal fibre connectivity. *Brain*, *135*(7), 2245-2255.

5

OPIOID USE IN INDIANA: CONSUMPTION PATTERNS AND CONSEQUENCES

INTRODUCTION

Opioids are a class of drugs that are used to reduce pain. They include legal substances such as prescription pain relievers received from a physician and illegal substances such as heroin or illicitly manufactured fentanyl. All opioids are chemically similar and the brain does not distinguish between legal and illegal opioids. By binding to special opioid receptors on nerve cells in the brain and body, opioids block pain signals and are responsible for the release of large amounts of dopamine. The release of dopamine has a strong reinforcing effect and is often experienced as “euphoria” and a “sense of wellbeing” in users (National Institute on Drug Abuse [NIDA], 2016, 2018a, 2018b).

Common prescription opioids include oxycodone (e.g., OxyContin®, Percocet®), hydrocodone (e.g., Vicodin®), oxymorphone (e.g., Opana®), codeine, morphine, and fentanyl (NIDA, 2018b). Fentanyl is a powerful synthetic opioid similar to morphine but 50 to 100 times stronger. The high potency of the drug significantly increases its risk for overdose. Fentanyl is typically used to treat severe pain or to manage pain after surgery. However, non-pharmaceutical fentanyl is sold on the streets in form of a powder, spiked on blotter


paper, and mixed with heroin or other drugs (NIDA, 2016). Prescription opioids are generally safe when taken for a short time and as prescribed by a healthcare provider. However, regular use, even as prescribed, can lead to dependence and addiction, and may result in overdose (NIDA, 2018b).

Heroin is a semi-synthetic illegal drug derived from morphine, a naturally occurring substance extracted from the opium poppy. Heroin is available in the form of a white or brown powder, or a black sticky substance commonly known as black tar heroin (NIDA, 2018a).

INSPECT

INSPECT is Indiana’s prescription drug monitoring program; it collects information on all controlled substances (DEA Schedules II through V) dispensed within the state. In 2018, nearly 5.7 million prescriptions for opioids were filled in Indiana, reflecting a dispensation rate of 214.3 per 1,000 residents. The number and rate of opioid dispensations has been gradually declining for the past two years (see Figure 5.1) (Indiana State Department of Health [ISDH], 2019). For county-level information see Map 5.1 and Appendix 5A.

Figure 5.1 Number and Rate (per 1,000 Population) of Opioids Dispensed in Indiana per Quarter (INSPECT, 2017-2018)



Note: Dispensation data includes three opioid prescription categories: (1) opioid analgesics, (2) opioid antidiarrheals/antitussives, and (3) opioid antagonists and treatment addiction medications.
Source: ISDH, 2019


**PREVALENCE OF OPIOID CONSUMPTION
IN THE GENERAL POPULATION**
National Survey on Drug Use and Health

Based on 2016–2017 averages from the Substance Abuse and Mental Health Services Administration (SAMHSA)’s National Survey on Drug Use and Health (NSDUH), an estimated 4.8% (95% Confidence Interval [CI]: 4.1–5.7) of Indiana residents ages 12 and older misused pain relievers in the past year (U.S.: 4.2%;

95% CI: 4.0–4.3). The highest rate was found among young adults ages 18 to 25, at 8.8% (95% CI: 7.1–10.9); this represents a rate statistically similar to the nation’s rate of 7.1% (95% CI: 6.8–7.5) (SAMHSA, 2017). For additional rates by age group, see Figure 5.2.

(In 2015, SAMHSA redesigned the questions in the NSDUH pertaining to the use and misuse of prescription drugs. Due to these changes, new estimates cannot be compared to previous years of the NSDUH.)

Figure 5.2 Prevalence of Past-Year Pain Reliever Use in Indiana and the United States, by Age Group (National Survey on Drug Use and Health, 2016-2017)




Source: SAMHSA, 2017

Although heroin use in the general U.S. population is relatively low (an estimated 0.3%), the percentage of Americans using the drug is higher than it was 10 years ago (Lipari and Hughes, 2015). Heroin has also become a major concern in Indiana. Its rise in use, as evidenced by the increase in heroin overdose fatalities, has led to several efforts by state agencies and organizations to identify and develop sources of Indiana-specific data and surveillance (Indiana State Department of Health [ISDH], 2019).

According to findings from the 2016–2017 NSDUH, 0.6% (95% CI: 0.4–1.1) of Hoosiers ages 12 and older reported using heroin in the past year; the U.S. rate was similar. Past-year heroin use was most prevalent among young adults ages 18 to 25, at 1.3% (95% CI: 0.7–2.3) (SAMHSA, 2017). For additional rates by age group, see Figure 5.3.

Figure 5.3 Percentage of Indiana and U.S. Population (12 years and older) Reporting Past-Year Heroin Use, by Age Group (National Survey on Drug Use and Health, 2016-2017)




Source: SAMHSA, 2017

Youth Risk Behavior Surveillance System

In 2015, 2.4% (95% CI: 1.3–4.4) of high school students (grades 9 through 12) in Indiana reported having tried heroin at least once in their life, according to the 2015 Youth Risk Behavior Surveillance System (YRBSS). Indiana’s rate was similar to the national rate of 2.1%

(95% CI: 1.5–2.8) (see Figure 5.4). No statistical differences by gender, race, or grade level were observed in 2015. Prevalence of lifetime heroin use has remained relatively stable among both Indiana and national high school students from 2005 through 2015 (Centers for Disease Control and Prevention [CDC], 1991–2015).

Figure 5.4 Percentage of Indiana and U.S. High School Students (Grades 9 through 12) Who Have Used Heroin at Least Once During their Lifetime (Youth Risk Behavior Surveillance System, 2003–2015)



Note: 2013 estimates for Indiana are not available.

Source: CDC, 1991–2015


As noted previously, a common route of administration for heroin is by needle injection. According to the 2015 YRBSS, the percentage of high school students who used a needle to inject any illegal drug into their body one or more times during their lifetime was statistically similar in Indiana (2.2%; 95% CI: 1.1–4.3) and the nation (1.8%; 95% CI: 1.3–2.3) (CDC, 1991–2015).

(While the YRBSS offers information on overall prescription drug misuse, it does not provide estimates for prescription pain reliever misuse specifically.)

Indiana Youth Survey


Based on results from the 2018 Indiana Youth Survey (INYS), past-month heroin use among 7th through 12th grade students was between 0.0% and 0.2% (see Figure 5.5). Heroin use among Indiana 12th graders peaked in 2011 at 1.2%, but is now at 0.2% (see Figure 5.6) (Gassman et al., 2018). For monthly heroin use rates in Indiana by region and grade level, see Appendix 5B.

Figure 5.5 Percentage of Indiana 7th through 12th Grade Students Reporting Monthly Heroin Use (Indiana Youth Survey, 2018)



Source: Gassman et al., 2018

Figure 5.6 Percentage of Indiana and U.S. 12th Grade Students Reporting Monthly Heroin Use (Indiana Youth Survey and Monitoring the Future Survey, 2009–2018)



Source: Gassman et al., 2018; Inter-university Consortium for Political and Social Research, University of Michigan, 2018

Indiana College Substance Use Survey

The Indiana College Substance Use Survey¹ includes questions on the past-month use of opioids and prescription painkillers not prescribed to the student.

Findings from the 2018 survey were as follows:

- a) Misuse of prescription painkillers:
 - 1.7% of Indiana college students misused prescription painkillers in the past month.
 - Rates did not differ significantly by gender.
 - Rates were higher among college students ages 21 to 25 (2.0%) compared to those under the age of 21 (1.5%).
- b) Misuse of heroin:
 - 0.2% of Indiana college students reported using heroin within the past month.
 - Rates did not differ significantly by gender or age group.

(King & Jun, 2018).

USE OF OPIOIDS IN THE TREATMENT POPULATION

Treatment Episode Data Set

Another method of tracking opioid misuse is to examine the Treatment Episode Data Set (TEDS) for individuals who report misuse of prescription pain relievers² or heroin at the time of substance use treatment admission.

In nearly 21% of Indiana treatment admissions, misuse of prescription opioids was reported (U.S.: 13.9%) and in over 10%, dependence³ was indicated in 2016 (SAMHSA, 2016). Significant differences in self-reported prescription opioid misuse were seen by gender, race, and age group. Generally speaking, women, whites, and adults between the ages of 25 and 44 had the highest percentages of misuse and dependence (see Table 5.1). Furthermore, the percentage of treatment admissions attributable to prescription opioids has increased from 2007 to 2016, with a peak in any prescription opioid misuse reported at admission in Indiana in 2014 (see Figure 5.7). For county-level information, see Appendix 5C.

Table 5.1 Percentage of Indiana and U.S. Treatment Episodes with Prescription Opioid Misuse and Dependence Reported at Treatment Admission, by Gender, Race, and Age Group (Treatment Episode Data Set, 2016)

		Misuse		Dependence	
		Indiana	U.S.	Indiana	U.S.
Gender	Male	17.6%	11.5%	8.5%	6.4%
	Female	25.9%	17.8%	13.1%	10.7%
Race	White	23.7%	17.1%	11.7%	9.7%
	Black	6.0%	5.9%	3.0%	3.3%
	Other	15.9%	8.9%	8.2%	4.7%
Age Group	Under 18	6.0%	5.0%	0.9%	1.2%
	18-24	17.8%	13.7%	6.9%	7.1%
	25-34	26.7%	18.2%	13.4%	10.5%
	35-44	22.4%	14.7%	12.0%	8.6%
	45-54	13.9%	9.0%	6.9%	5.2%
	55+	10.6%	8.6%	7.5%	5.2%
Total		20.9%	13.9%	10.4%	7.9%


Source: SAMHSA, 2016

¹Twenty-four (24) colleges participated in the 2018 survey; results are based on nonrandom sampling and are not representative of all college students in Indiana.

²We used TEDS variables “nonprescription methadone” and “other opiates/synthetics” to define pain reliever use (excludes heroin).

³We defined prescription pain reliever dependence as “individuals in substance abuse treatment listing prescription pain relievers as their primary substance at admission.”

Figure 5.7 Percentage of Indiana and U.S. Treatment Episodes with Prescription Opioid Misuse and Dependence Reported at Treatment Admission (Treatment Episode Data Set, 2007–2016)




Source: SAMHSA, 2016

In over one-fifth of Indiana treatment admissions in 2016, heroin use was reported; heroin dependence⁴ was indicated in 17.5% of admissions (SAMHSA, 2016). Though the percentage of treatment admissions

attributable to heroin in Indiana increased significantly from 2007 through 2016, Indiana’s percentage remained below the U.S. percentage. For additional trend information, see Figure 5.8.

Figure 5.8 Percentage of Indiana and U.S. Treatment Episodes with Heroin Use and Dependence Reported at Treatment Admission (Treatment Episode Data Set, 2007–2016)



Source: SAMHSA, 2016

⁴We defined heroin dependence as “individuals in substance abuse treatment listing heroin as their primary substance at admission.”

Within Indiana’s treatment population, reported heroin use and dependence differed significantly by gender, race, and age group:

- **Gender**—Reported heroin use and dependence increased among both males and females, but has consistently been higher in females.
- **Race**—Whites had the highest percentage of heroin use and dependence compared to all other races.

- **Age**—Heroin use and dependence within Indiana’s treatment population was highest among adults ages 25 to 34.

For additional details, see Table 5.2 (SAMHSA, 2016); for county-level information, see Appendix 5C.

Table 5.2 Percentage of Indiana and U.S. Treatment Episodes with Heroin Use and Dependence Reported at Treatment Admission, by Gender, Race, and Age Group (Treatment Episode Data Set, 2016)

		Use		Dependence	
		Indiana	U.S.	Indiana	U.S.
Gender	Male	20.0%	30.0%	15.6%	25.9%
	Female	26.1%	31.5%	20.4%	27.2%
Race	White	24.7%	33.2%	19.2%	28.6%
	Black	8.5%	21.9%	7.3%	18.6%
	Other	22.0%	29.1%	17.4%	25.0%
Age Group	Under 18	1.7%	2.7%	1.1%	1.7%
	18-24	23.8%	32.5%	18.7%	28.4%
	25-34	31.3%	38.0%	24.9%	33.1%
	35-44	19.9%	29.1%	14.6%	24.7%
	45-54	10.0%	25.3%	7.5%	21.2%
	55+	8.5%	24.3%	6.5%	21.0%
Total		22.4%	30.6%	17.5%	26.3%

Source: SAMHSA, 2016

Opioid Treatment Programs

Opioid treatment programs (OTPs) provide medication-assisted treatment to individuals with an opioid use disorder. OTPs are certified by SAMHSA, accredited by an independent SAMHSA-approved accrediting body, and licensed by the state in which they operate. Federal law requires OTPs to provide medical, counseling, vocational, educational, and other assessment and treatment services, in addition to prescribed medication. In 2017, a total of 13,697 unique patients were treated in OTPs in Indiana (Indiana Family and Social Services Administration, 2018b).


CONSEQUENCES OF OPIOID USE Fatal and Non-Fatal Drug Overdoses

In high doses and/or combined with alcohol and certain other drugs, opioids can cause respiratory depression and lead to death (NIDA, 2018a). Drug overdose deaths (from all drugs) increased in Indiana from 9.7 per 100,000 population (U.S.: 10.1) in 2005 to 29.4 per 100,000 population (U.S.: 21.7) in 2017 (CDC, 1999–2017).⁵ A large percentage of overall drug overdoses involve opioids. In Indiana, the number of overdose deaths involving an opioid rose from 347 in 2011 to 1,138 in 2017 (ISDH, 2019). For 2011 through 2017 overdose mortality rates involving opioids, see Figure 5.9.

In addition, a total of 8,169 visits to Indiana emergency departments occurred due to a nonfatal opioid overdose in 2017 (ISDH, 2019).

⁵Includes ICD-10 causes of death: X40-X44, X60-X64, X85, Y10-Y14.

Figure 5.9 Drug Overdose Deaths Involving Opioids, Rate per 100,000 Population (Indiana, 2011–2017)



Note: “Rx (prescription) Opioid” and “Heroin” are subcategories of “Any Opioid”. Overdose deaths involving prescription opioids or heroin are not mutually exclusive as multiple drugs are frequently involved in overdose deaths. Source: ISDH, 2019

HIV/AIDS and Hepatitis B & C

Opioids, especially when injected, are a significant risk factor for contracting human immunodeficiency virus infection (HIV) and hepatitis B and C, due to the common practice of needle-sharing among injection drug users (NIDA, 2018c). However, drug use in any form is associated with risk behaviors related to infectious disease transmission (NIDA, 2018c).

As of December 31, 2017, a total of 12,635 individuals in Indiana were living with HIV or AIDS, representing an annual HIV/AIDS prevalence rate of 190.5 per 100,000 population. In 2017, there were 547 new cases of HIV/AIDS. In nearly 9% of these new cases, injection drug use (IDU) was reported, either as the sole risk factor for contracting HIV/AIDS or in combination with other risk factors (ISDH, 2019).

Indiana’s age-adjusted HIV/AIDS mortality rate for 2016 was 1.0 per 100,000 population (95% CI: 0.8–1.3), which was similar to the U.S. rate of 1.6 per 100,000 population (95% CI: 1.6–1.6) (CDC, 1999–2017).⁶

Hepatitis is a liver disease that is caused by viral infection. The hepatitis B virus (HBV) and hepatitis C virus (HCV) are transmitted when blood of an infected person enters the body of a person who is not infected. Injection drug use (IDU) is a major risk factor for both acquiring and transmitting HBV and HCV. It is estimated that each injection drug user infected with HCV is likely to infect 20 other people, extending the risk of infection far beyond the individual using the drug (NIDA, 2018d).

In 2017, there were 905 cases of hepatitis B (including 170 acute and 735 chronic cases) and 8,222 cases of hepatitis C (including 236 acute and 7,986 chronic cases) in Indiana (ISDH, 2019).

The 2017 age-adjusted mortality rate attributable to HBV and HCV⁷ combined was 1.0 per 100,000 population (95% CI: 0.8–1.2) in Indiana, which was below the national rate of 1.3 per 100,000 population (95% CI: 1.3–1.4) (CDC, 1999–2017).

⁶Mortality rates for HIV/AIDS are based on ICD-10 codes B20-B24 (Human immunodeficiency virus [HIV] disease).

⁷Mortality rates for hepatitis B and C infections are based on the following ICD-10 codes: B16 (Acute hepatitis B), B17.0 (Acute delta-[super]infection of hepatitis B carrier), B17.1 (Acute hepatitis C), B18.0 (Chronic viral hepatitis B with delta-agent), B18.1 (Chronic viral hepatitis B without delta-agent), B18.2 (Chronic viral hepatitis C).

Pharmacy Robberies

The number of pharmacy robberies in Indiana peaked in 2015 with 175 robberies, but decreased to 22 robberies in 2017. These 22 robberies were responsible

for nearly 37,000 doses of stolen controlled substances, reflecting a purchase value of \$76,439 (see Table 5.3) (IPLA, 2018).

Table 5.3 Pharmacy Robberies in Indiana (Summary Report)

	2013	2014	2015	2016	2017
Number of Robberies	71	80	175	75	22
Purchase Value of Stolen Drugs	\$202,133	\$293,079	\$479,785	\$246,138	\$76,439
Number of Doses Stolen					
Oxycodone	6,457	50,525	113,807	46,325	30,911
Hydrocodone	4,159	14,702	40,452	25,737	-
Others	1,735	10,502	29,442	34,859	6,028
Total	12,351	75,729	183,701	106,921	36,939

Source: IPLA, 2018

APPENDIX 5A

Number and Rate (per 1,000 Population) of Opioid Dispensations in Indiana, by County of Patient's Residence (INSPECT, 2018)

County	Number of Dispensations	Rate of Dispensations per 1,000
Adams	20,392	144.7
Allen	257,490	173.8
Bartholomew	76,147	233.9
Benton	6,665	192.6
Blackford	15,452	318.0
Boone	47,782	184.8
Brown	15,957	267.5
Carroll	15,651	195.9
Cass	31,498	207.5
Clark	125,301	270.0
Clay	23,381	222.2
Clinton	30,481	234.8
Crawford	12,596	298.8
Daviess	28,819	218.5
Dearborn	48,381	245.2
Decatur	25,875	243.2
DeKalb	33,856	198.0
Delaware	122,018	263.9
Dubois	33,159	194.8
Elkhart	125,697	154.2
Fayette	35,520	380.6
Floyd	75,511	245.2
Fountain	15,321	232.3
Franklin	20,518	225.8
Fulton	19,324	239.9
Gibson	35,365	262.3
Grant	76,943	287.4
Greene	36,111	280.3
Hamilton	170,673	134.9
Hancock	63,497	215.3
Harrison	40,301	253.0
Hendricks	111,973	174.3
Henry	64,698	333.4
Howard	99,454	301.1
Huntington	33,601	230.8
Jackson	40,716	231.3
Jasper	35,934	268.7
Jay	18,675	221.8
Jefferson	39,569	305.1
Jennings	31,724	285.7
Johnson	133,866	220.2
Knox	50,221	332.6
Kosciusko	58,397	184.6
LaGrange	16,762	107.1
Lake	380,600	195.8
LaPorte	113,924	258.9
Lawrence	65,122	357.7

County	Number of Dispensations	Rate of Dispensations per 1,000
Madison	158,760	307.0
Marion	699,407	185.8
Marshall	38,125	204.7
Martin	13,142	323.0
Miami	32,633	227.4
Monroe	96,602	166.0
Montgomery	37,440	245.8
Morgan	81,748	293.2
Newton	13,643	245.0
Noble	39,328	206.4
Ohio	6,545	275.8
Orange	21,934	283.6
Owen	27,367	328.3
Parke	12,164	181.0
Perry	16,239	214.1
Pike	16,594	333.7
Porter	159,255	237.3
Posey	28,503	279.7
Pulaski	14,466	285.7
Putnam	32,763	218.8
Randolph	25,785	257.0
Ripley	24,949	216.2
Rush	17,477	262.4
St. Joseph	193,999	180.2
Scott	32,693	344.4
Shelby	42,201	238.0
Spencer	16,910	204.7
Starke	32,829	356.7
Steuben	24,001	175.9
Sullivan	19,322	232.2
Switzerland	9,776	232.2
Tippecanoe	111,230	147.9
Tipton	15,945	262.6
Union	4,627	160.4
Vanderburgh	199,927	275.0
Vermillion	14,869	237.6
Vigo	91,132	211.1
Wabash	35,578	280.0
Warren	6,245	191.2
Warrick	57,868	231.5
Washington	33,426	302.0
Wayne	75,770	284.6
Wells	22,827	204.2
White	21,026	219.0
Whitley	28,745	214.8
Indiana	5,687,064	214.3

Note: Dispensation data includes three opioid prescription categories: (1) opioid analgesics, (2) opioid antidiarrheals/antitussives, and (3) opioid antagonists and treatment addiction medications.

Source: ISDH, 2019

APPENDIX 5B

Percentage of Indiana Students Reporting Monthly Heroin Use, by Region and Grade (Indiana Youth Survey, 2018)

	Indiana	Northwest	North Central	Northeast	West	Central	East	Southwest	Southeast
7th Grade	0.0	0.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0
8th Grade	0.1	0.1	0.1	0.3	0.1	0.0	0.1	0.1	0.1
9th Grade	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.1
10th Grade	0.1	0.2	0.0	0.0	0.1	0.1	0.1	0.5*	0.1
11th Grade	0.2	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.2
12th Grade	0.2	0.3	0.3	0.0	0.2	0.1	0.2	0.1	0.0

Notes: * Indicates a local rate that is significantly different from the overall state rate ($P < 0.05$).

Beginning in 2015, the Indiana Youth Survey stopped asking 6th grade students about heroin use.

Source: Gassman et al., 2018

APPENDIX 5C

Number of Treatment Episodes with Prescription (Rx) Opioid Misuse and Dependence and Heroin Use and Dependence Reported at Treatment Admission in Indiana, by County (Substance Abuse Population by County/ Treatment Episode Data Set, 2018)

County	Treatment Episodes	Rx Opioid Misuse		Rx Opioid Dependence		Heroin Use		Heroin Dependence	
	Total	Number	%	Number	%	Number	%	Number	%
Adams	228	46	20.2%	17	7.5%	37	16.2%	21	9.2%
Allen	2,050	324	15.8%	160	7.8%	334	16.3%	254	12.4%
Bartholomew	727	139	19.1%	75	10.3%	224	30.8%	164	22.6%
Benton	61	9	14.8%	<5	N/A	6	9.8%	<5	N/A
Blackford	144	31	21.5%	14	9.7%	77	53.5%	63	43.8%
Boone	167	51	30.5%	17	10.2%	61	36.5%	43	25.7%
Brown	72	12	16.7%	9	12.5%	26	36.1%	18	25.0%
Carroll	80	13	16.3%	8	10.0%	10	12.5%	5	6.3%
Cass	387	49	12.7%	17	4.4%	51	13.2%	32	8.3%
Clark	103	44	42.7%	35	34.0%	21	20.4%	13	12.6%
Clay	106	13	12.3%	7	6.6%	6	5.7%	<5	N/A
Clinton	147	39	26.5%	22	15.0%	34	23.1%	21	14.3%
Crawford	36	8	22.2%	5	13.9%	<5	N/A	<5	N/A
Daviess	202	49	24.3%	26	12.9%	21	10.4%	10	5.0%
Dearborn	497	195	39.2%	86	17.3%	161	32.4%	108	21.7%
Decatur	185	42	22.7%	20	10.8%	24	13.0%	16	8.6%
DeKalb	276	48	17.4%	26	9.4%	21	7.6%	14	5.1%
Delaware	843	287	34.0%	161	19.1%	291	34.5%	231	27.4%
Dubois	131	35	26.7%	18	13.7%	7	5.3%	<5	N/A
Elkhart	765	93	12.2%	57	7.5%	68	8.9%	45	5.9%
Fayette	322	121	37.6%	69	21.4%	132	41.0%	96	29.8%
Floyd	49	22	44.9%	16	32.7%	13	26.5%	10	20.4%
Fountain	59	13	22.0%	8	13.6%	6	10.2%	<5	N/A
Franklin	145	49	33.8%	28	19.3%	42	29.0%	29	20.0%
Fulton	188	20	10.6%	6	3.2%	35	18.6%	20	10.6%
Gibson	252	47	18.7%	19	7.5%	6	2.4%	<5	N/A
Grant	509	106	20.8%	46	9.0%	203	39.9%	149	29.3%
Greene	197	50	25.4%	28	14.2%	25	12.7%	12	6.1%
Hamilton	874	141	16.1%	50	5.7%	166	19.0%	126	14.4%
Hancock	398	62	15.6%	29	7.3%	83	20.9%	70	17.6%
Harrison	19	<5	N/A	<5	N/A	9	47.4%	8	42.1%
Hendricks	424	58	13.7%	22	5.2%	108	25.5%	85	20.0%
Henry	381	151	39.6%	104	27.3%	55	14.4%	34	8.9%
Howard	492	120	24.4%	29	5.9%	183	37.2%	108	22.0%
Huntington	136	47	34.6%	15	11.0%	43	31.6%	28	20.6%
Jackson	330	90	27.3%	30	9.1%	73	22.1%	36	10.9%
Jasper	93	22	23.7%	5	5.4%	30	32.3%	21	22.6%
Jay	122	22	18.0%	7	5.7%	67	54.9%	54	44.3%
Jefferson	307	108	35.2%	52	16.9%	40	13.0%	24	7.8%
Jennings	260	44	16.9%	15	5.8%	39	15.0%	24	9.2%
Johnson	424	79	18.6%	47	11.1%	128	30.2%	97	22.9%
Knox	352	86	24.4%	55	15.6%	12	3.4%	<5	N/A

(continued on next page)

APPENDIX 5C (Continued from previous page)

County	Treatment Episodes	Rx Opioid Misuse		Rx Opioid Dependence		Heroin Use		Heroin Dependence	
	Total	Number	%	Number	%	Number	%	Number	%
Kosciusko	362	73	20.2%	26	7.2%	76	21.0%	47	13.0%
LaGrange	115	19	16.5%	11	9.6%	<5	N/A	<5	N/A
Lake	1,903	199	10.5%	115	6.0%	470	24.7%	408	21.4%
LaPorte	172	31	18.0%	19	11.0%	73	42.4%	61	35.5%
Lawrence	535	208	38.9%	88	16.4%	113	21.1%	77	14.4%
Madison	1,205	425	35.3%	202	16.8%	222	18.4%	120	10.0%
Marion	4,134	554	13.4%	314	7.6%	1000	24.2%	867	21.0%
Marshall	137	26	19.0%	12	8.8%	25	18.2%	17	12.4%
Martin	60	12	20.0%	6	10.0%	<5	N/A	<5	N/A
Miami	249	46	18.5%	14	5.6%	64	25.7%	35	14.1%
Monroe	1,428	388	27.2%	194	13.6%	357	25.0%	215	15.1%
Montgomery	304	52	17.1%	17	5.6%	81	26.6%	57	18.8%
Morgan	550	123	22.4%	43	7.8%	126	22.9%	93	16.9%
Newton	27	8	29.6%	<5	N/A	11	40.7%	9	33.3%
Noble	346	35	10.1%	17	4.9%	18	5.2%	7	2.0%
Ohio	33	8	24.2%	<5	N/A	9	27.3%	6	18.2%
Orange	135	25	18.5%	18	13.3%	9	6.7%	5	3.7%
Owen	187	32	17.1%	17	9.1%	28	15.0%	15	8.0%
Parke	65	6	9.2%	<5	N/A	11	16.9%	<5	N/A
Perry	94	14	14.9%	6	6.4%	<5	N/A	<5	N/A
Pike	46	9	19.6%	5	10.9%	<5	N/A	<5	N/A
Porter	202	50	24.8%	26	12.9%	82	40.6%	70	34.7%
Posey	156	33	21.2%	13	8.3%	<5	N/A	<5	N/A
Pulaski	82	17	20.7%	11	13.4%	9	11.0%	7	8.5%
Putnam	253	36	14.2%	15	5.9%	22	8.7%	12	4.7%
Randolph	167	38	22.8%	24	14.4%	68	40.7%	58	34.7%
Ripley	180	53	29.4%	28	15.6%	40	22.2%	29	16.1%
Rush	139	38	27.3%	18	12.9%	16	11.5%	10	7.2%
Saint Joseph	1,351	141	10.4%	60	4.4%	325	24.1%	264	19.5%
Scott	212	68	32.1%	39	18.4%	50	23.6%	34	16.0%
Shelby	140	36	25.7%	13	9.3%	43	30.7%	28	20.0%
Spencer	132	22	16.7%	11	8.3%	5	3.8%	<5	N/A
Starke	77	30	39.0%	16	20.8%	26	33.8%	21	27.3%
Steuben	183	21	11.5%	5	2.7%	19	10.4%	13	7.1%
Sullivan	115	34	29.6%	19	16.5%	7	6.1%	<5	N/A
Switzerland	113	39	34.5%	16	14.2%	29	25.7%	16	14.2%
Tippecanoe	364	63	17.3%	22	6.0%	83	22.8%	63	17.3%
Tipton	46	9	19.6%	5	10.9%	<5	N/A	<5	N/A
Union	54	22	40.7%	<5	N/A	20	37.0%	18	33.3%
Vanderburgh	1,218	264	21.7%	118	9.7%	99	8.1%	64	5.3%
Vermillion	132	20	15.2%	11	8.3%	18	13.6%	7	5.3%
Vigo	771	79	10.2%	34	4.4%	40	5.2%	25	3.2%
Wabash	299	99	33.1%	34	11.4%	95	31.8%	64	21.4%
Warren	26	5	19.2%	<5	N/A	<5	N/A	<5	N/A

(continued on next page)

APPENDIX 5C (Continued from previous page)

County	Treatment Episodes	Rx Opioid Misuse		Rx Opioid Dependence		Heroin Use		Heroin Dependence	
	Total	Number	%	Number	%	Number	%	Number	%
Warrick	310	59	19.0%	29	9.4%	15	4.8%	11	3.5%
Washington	44	16	36.4%	8	18.2%	15	34.1%	14	31.8%
Wayne	558	129	23.1%	61	10.9%	234	41.9%	193	34.6%
Wells	156	55	35.3%	21	13.5%	42	26.9%	31	19.9%
White	125	16	12.8%	<5	N/A	9	7.2%	<5	N/A
Whitley	126	36	28.6%	16	12.7%	19	15.1%	10	7.9%
County Info Missing	41	9	22.0%	<5	N/A	13	31.7%	11	26.8%
Indiana	33,399	6,815	20.4%	3,257	9.8%	7,114	21.3%	5,201	15.6%

Notes: We defined prescription opioid dependence as “individuals in substance abuse treatment listing prescription opioids as their primary substance at admission.”

We defined heroin dependence as “individuals in substance abuse treatment listing heroin as their primary substance at admission.”

We calculated the percentages by dividing the number of reported prescription drug use/dependence by the number of treatment episodes.

Information on treatment episodes <5 was suppressed due to confidentiality constraints.

Source: Indiana Family and Social Services Administration, 2018a

REFERENCES, CHAPTER 5

- Centers for Disease Control and Prevention. (1991-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (1999-2017). *CDC WONDER underlying causes of death (compressed mortality)*. Retrieved from <http://wonder.cdc.gov/>
- Gassman, R., Jun, M., Samuel, S., Agle, Lee, J., & Wolf, J. (2018). *Indiana Youth Survey*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Indiana Family and Social Services Administration. (2018a). *Treatment Episode Data System (TEDS)*. Indianapolis, IN: Indiana Family and Social Services Administration.
- Indiana Family and Social Services Administration. (2018b). *Opioid Treatment Programs – Patient Count*. Email correspondence with Rhonda Webb from February 15, 2018.
- Indiana Professional Licensing Agency. (2018). *Pharmacy robberies in Indiana 2017 – summary report*. Email correspondence with Zaneta Nunnally from March 28, 2018.
- Indiana State Department of Health. (2019). *Stats Explorer*. Retrieved from https://gis.in.gov/apps/isdh/meta/stats_layers.htm
- Inter-university Consortium for Political and Social Research, University of Michigan. (2018). *Monitoring the Future (MTF)*. Retrieved from <http://www.monitoringthefuture.org/data/data.html>
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey, 2018*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://drugs.indiana.edu/indiana-college-survey/substance-use-survey>
- National Institute on Drug Abuse. (2016). *DrugFacts: Fentanyl*. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/fentanyl>
- National Institute on Drug Abuse. (2018a). *DrugFacts: Heroin*. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/heroin>
- National Institute on Drug Abuse. (2018b). *DrugFacts: Prescription Opioids*. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/prescription-opioids>
- National Institute on Drug Abuse. (2018c). *Drug Use and Viral Infections (HIV, Hepatitis)*. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/drug-use-viral-infections-hiv-hepatitis>
- National Institute on Drug Abuse. (2018d). *Why does heroin use create special risk for contracting HIV/AIDS and hepatitis B and C?* Retrieved from <https://www.drugabuse.gov/publications/research-reports/heroin/why-are-heroin-users-special-risk-contracting-hiv-aids-hepatitis-b-c>
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2016). *Treatment Episode Data Set -- Admissions (TEDS-A)*. Retrieved from <http://www.dasis.samhsa.gov/dasis2/teds.htm>
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>

STIMULANT USE IN INDIANA: CONSUMPTION PATTERNS AND CONSEQUENCES

INTRODUCTION

Stimulants encompass a group of both legal and illicit drugs that share similar physiological mechanisms of action. When ingested, stimulants lead to an increase in alertness, attention, and energy while also elevating blood pressure, heart rate, and respiration. In the brain, stimulants raise dopamine levels which can lead to feelings ranging from pleasure to intense euphoria. Stimulant use is also often associated with feelings of increased wakefulness, motivation, mental focus, and libido (National Institute on Drug Abuse [NIDA], 2018). While a number of stimulant drugs exist, the three associated with the greatest level of problematic use are cocaine, methamphetamine, and prescription stimulants.

Cocaine is a highly addictive stimulant produced from the leaves of the coca plant. The two most common forms of cocaine are powder cocaine and crack cocaine. Powder cocaine is a fine white powder and, while it can be injected, is most often snorted or inhaled. Crack cocaine is cocaine that has been processed into a rock crystal. Crack is typically used by placing the crystals into a glass pipe, heating them, and then inhaling the vapors. The name “crack” refers to the crackling sound made when the rock is heated (NIDA, 2016a, 2016b). Both forms of cocaine increase levels of dopamine in the brain resulting in a short-lived, intense high that can range from 15 to 30 minutes for powder cocaine or 5 to 10 minutes for crack cocaine.

Methamphetamine (meth), also known as “crystal” or “ice”, is a highly addictive stimulant derived from amphetamine. Although meth can be taken in a variety of ways, most users in Indiana report either smoking it or injecting it intravenously (Indiana Family and Social Services Administration [FSSA], 2017; NIDA, 2017). Upon initial administration, meth users experience a short, intense euphoria or “rush” followed by an extended high that can last up to 12 hours due to

the drug’s long half-life (Halkitis, Parsons, & Stirrat, 2001; Centers for Disease Control and Prevention [CDC], 2007). The intensity of meth stimulation depends on the mode of administration. Oral ingestion or snorting produces a longer-lasting, but less intense effect, while smoking or injecting results in a briefer but more intense rush (Homer et al., 2008).


Prescription stimulants are legally produced stimulants such as dextroamphetamine (Dexedrine), methylphenidate (Ritalin), amphetamine sulfate (Adderall), and lisdexamfetamine (Vyvanse). These drugs increase alertness, attention, and energy and are used for the treatment of narcolepsy and attention-deficit hyperactivity disorder. Although some people may choose to use prescription stimulants as a way to get high, many individuals who use these drugs inappropriately may do so in an attempt to enhance academic/work performance or improve memory (NIDA, 2018).

PREVALENCE OF STIMULANT CONSUMPTION IN THE GENERAL POPULATION

National Survey on Drug Use and Health


The National Survey on Drug Use and Health (NSDUH) estimated that in 2017, approximately 1.9% (95% Confidence Interval [CI]: 1.4–2.5) of Hoosiers 12 years of age or older used cocaine in the past year, an estimate similar to that for the nation (2.0%; 95% CI: 1.9–2.1). Across age groups, cocaine use was highest among persons between the ages of 18 and 25 in both Indiana (5.7%, 95% CI: 4.2–7.8) and the U.S. (5.9%, 95% CI: 5.5–6.3) (see Figure 6.1). Over the last 10 years, the rate of past-year cocaine use in both Indiana and the U.S. has remained fairly stable (see Figure 6.2) (Substance Abuse and Mental Health Services Administration [SAMHSA], 2017).

Figure 6.1 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Cocaine Use in the Past Year, by Age Group (National Survey on Drug Use and Health, 2017)



Source: SAMHSA, 2017

Figure 6.2 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Cocaine Use in the Past Year (National Survey on Drug Use and Health, 2008-2017)



Source: SAMHSA, 2017

2017 is the first year in which state-level NSDUH estimates on methamphetamine use were available. In Indiana, 0.6% of Hoosiers (95% CI: 0.4-1.0) reported using meth in the past year; the U.S. rate was the same (0.6%; 95% CI: 0.5-0.6). For prevalence rates by age group, see Figure 6.3 (SAMHSA, 2017).

Youth Risk Behavior Surveillance Survey

According to the 2015 Youth Risk Behavior Surveillance System (YRBSS), 4.0% (95% CI: 2.9–5.7) of Indiana high school students (grades 9-12) reported that they had used a form of cocaine at least once in their lifetime.


National rates for lifetime use were similar, at 5.2% (95% CI: 4.3–6.2). The difference in Indiana prevalence rates by gender, race/ethnicity, or grade level was not statically significant (see Table 6.1) (CDC, 1991-2015). The YRBSS estimated that in 2015, 2.9% (95% CI: 1.5–5.4) of Indiana high school students and a similar percentage of U.S. high school students (3.0%; 95% CI: 2.4–3.8) had ever used meth. Since 2003, the percentage of Indiana’s high school students estimated to have used either cocaine or meth has gradually declined (see Figure 6.4). The YRBSS does not ask students to describe their use of prescription stimulants.

Table 6.1 Percentage of Indiana and U.S. High School Students (Grades 9 through 12) Reporting Lifetime Cocaine or Methamphetamine Use, by Gender, Race/Ethnicity, and Grade (Youth Risk Behavior Surveillance System, 2015)

		Cocaine		Methamphetamine	
		Indiana (95%)	U.S. (95%)	Indiana	U.S.
Gender	Male	5.2% (3.4–7.9)	6.3% (5.1–7.9)	4.1% (2.0–8.2)	3.6% (2.6–4.9)
	Female	2.7% (1.7–4.2)	3.8% (3.1–4.6)	1.4% (0.8–2.6)	2.3% (1.7–3.0)
Race/Ethnicity	White	3.6% (2.3–5.6)	4.1% (3.3–5.2)	2.4% (1.1–5.3)	2.1% (1.5–2.8)
	Black	3.7% (1.2–10.7)	3.8% (2.5–6.0)	3.7% (1.2–10.7)	2.8% (1.5–5.1)
	Hispanic	7.9% (4.2–14.1)	8.0% (6.6–9.7)	3.2% (1.4–7.0)	4.4% (3.3–5.9)
Grade	9	3.5% (1.6–7.2)	3.4% (2.6–4.5)	3.5% (1.6–7.8)	2.0% (1.5–2.7)
	10	4.7% (3.4–6.5)	5.1% (3.8–6.8)	2.3% (1.4–3.8)	3.3% (2.3–4.9)
	11	4.7% (2.6–8.6)	5.0% (3.9–6.5)	3.7% (1.5–8.9)	2.8% (1.9–4.0)
	12	3.4% (1.8–6.3)	7.2% (5.6–9.1)	1.6% (0.4–6.6)	3.8% (2.7–5.3)
Total		4.0% (2.9–5.7)	5.2% (4.3–6.2)	2.9% (1.5–5.4)	3.0% (2.4–3.8)

Source: SAMHSA, 2014

Figure 6.3 Percentage of Indiana and U.S. Population (12 Years and Older) Reporting Methamphetamine Use in the Past Year, by Age Group (National Survey on Drug Use and Health, 2017)




Source: SAMHSA, 2017

Indiana Youth Survey and Monitoring the Future Survey

Both the Indiana Youth Survey (INYS) and the Monitoring the Future survey (MTF) provide local and national estimates, respectively, of current cocaine and methamphetamine use among 8th, 10th, and 12th grade students. Neither survey asks students to report on their current inappropriate use of prescription stimulants. According to the 2018 INYS, only a small percentage of Indiana’s 8th, 10th, and 12th graders reported currently


using either cocaine or meth. Current use of both substances has been decreasing in Indiana over the past 10 years and these decreases are consistent with national trends (see Figures 6.5 and 6.6) (Gassman et al., 2018; Inter-university Consortium for Political and Social Research [ICPSR], 2018). For 2018 data on current cocaine/crack use and meth use among students in grades 7 through 12 by Indiana region, see Appendix 6A.

Figure 6.4 Percentage of Indiana and U.S. High School Students (9th-12th Grade) Reporting Lifetime Methamphetamine Use (Youth Risk Behavior Surveillance System, 2003-2015)




Note: 2013 YRBSS data are not available for Indiana due to insufficient response rate.
Source: CDC, 1991-2015

Figure 6.5 Percentage of 8th, 10th, and 12th Grade Students Reporting Current Cocaine/Crack Use (Indiana Youth Survey and Monitoring the Future Survey, 2009-2018)



Source: Gassman et al., 2018; ICPSR, 2018

Figure 6.6 Percentage of 8th, 10th, and 12th Grade Students Reporting Current Meth Use (Indiana Youth Survey and Monitoring the Future Survey, 2009-2018)



Source: Gassman et al., 2018; ICPSR, 2018

The Indiana College Substance Use Survey

The Indiana College Substance Use Survey (ICSUS) provides estimates of alcohol, tobacco, and other drug use among Indiana college students. According to findings from the 2018 survey, which were based on 24 participating colleges and universities:

- 2.9% of Indiana college students reported having used cocaine in the past month,
- 0.2% reported having used meth, and
- 5.3% reported having used prescription stimulants not prescribed to them.

Most students who had used these drugs reported initiating use after entering college (74.9% cocaine; 68.4% methamphetamine; 54.8% prescription stimulants). Cocaine, meth, and prescription stimulants were used more frequently by students who were 21 years of age or older compared to those under 21 (King & Jun, 2018)¹.

USE OF STIMULANTS IN THE TREATMENT POPULATION

Treatment Episode Data Set

Data from the Treatment Episode Data Set (TEDS) indicate that methamphetamine was the most widely used stimulant in Indiana's substance use treatment population. In nearly one-fourth (23.7%) of treatment

admissions in Indiana methamphetamine use was reported in 2016; this represents a significantly higher proportion compared to the rest of the country (16.5%, $P < .001$). Methamphetamine use was more commonly reported among women, whites, and younger adults (see Table 6.2). The use of methamphetamine in Indiana's treatment population increased by more than 150% since 2007 (see Figure 6.7).

Cocaine was the second most frequently used stimulant in Indiana's treatment population and reported in 12.1% of treatment admissions in 2016; a percentage significantly lower than that noted for the rest of the nation (17.7%, $P < .001$). Cocaine use was reported more often by women, black individuals, and persons 45 years of age and older (see Table 6.2). The use of cocaine among those in treatment dropped nearly 50% since 2007 (see Figure 6.8).

Misuse of prescription stimulants was comparatively low within this population. In 1.7% of Indiana treatment admissions, misuse of these drugs was reported in 2016; a percentage similar to that for the rest of the country (1.6%). Aside from a spike in 2011-2012, the misuse of prescription stimulants by Indiana's treatment population has changed little over the past 10 years (see Figure 6.9). Women and whites entering treatment were more likely to report misuse (see Table 6.2).


¹Twenty-four colleges participated in the 2018 survey; results are based on nonrandom sampling and are not representative of all college students in Indiana.

Table 6.2 Stimulant Misuse Reported at Substance Use Treatment Admission in Indiana (Treatment Episode Data Set, 2016)

		Methamphetamine		Cocaine		Prescription Stimulants	
		N	%	N	%	N	%
Gender	Male	2,556	20.1%	1,463	11.5%	176	1.4%
	Female	2,492	28.9%	1,125	13.1%	196	2.3%
Race	White	4,771	27.5%	1,634	9.4%	337	1.9%
	Black	67	2.4%	781	27.9%	18	0.6%
	Other	209	17.6%	173	14.6%	17	1.4%
Ethnicity	Hispanic	186	12.7%	210	14.4%	27	1.8%
	Non-Hispanic	3,589	18.0%	2,162	10.9%	328	1.6%
Age	Under 18	30	4.6%	16	2.5%	13	2.0%
	18 to 24	905	23.0%	239	6.1%	73	1.9%
	25 to 34	2,330	28.5%	832	10.2%	161	2.0%
	35 to 44	1,245	26.9%	681	14.7%	82	1.8%
	45 to 54	435	16.5%	546	20.7%	32	1.2%
	55 or Older	103	8.0%	274	21.3%	11	0.9%
Total		5,048	23.7%	2,588	12.1%	372	1.7%


Source: SAMHSA, 2016

Figure 6.7 Percentage of Treatment Episodes with Reported Meth Use and Dependence, Indiana and the United States (Treatment Episode Data Set, 2007-2016)



Source: SAMHSA, 2016

Figure 6.8 Percentage of Treatment Episodes with Reported Cocaine Use and Dependence, Indiana and the United States (Treatment Episode Data Set, 2007-2016)



Source: SAMHSA, 2016

HEALTH CONSEQUENCES

The use of cocaine, meth, and prescription stimulants can all result in serious health consequences if used at high doses, especially over long periods of time. Ingesting large amounts of any of these drugs can result in serious cardiovascular, nervous system, or gastrointestinal complications, overdose, and in severe cases, death. Consuming stimulants can also lead to psychotic-like symptoms and paranoia, which, depending on the drug used, can be permanent. Meth use is particularly damaging to the body with long-term use associated with brain, liver, and kidney damage and serious dental problems (i.e., meth mouth). Although stimulant users who inject place themselves at particularly high risk for contracting blood-borne illnesses such as HIV and hepatitis, all stimulant users are at heightened risk for these illnesses as these drugs can severely impair judgment and lead to risky sexual behaviors with infected partners (NIDA, 2016a, 2017, 2018).


LEGAL CONSEQUENCES

Indiana State Police Meth Lab Seizures

Much of the meth currently consumed in the U.S. is produced in “superlabs,” most of which are located in Mexico (NIDA, 2017). However, because meth can


be produced using easily accessible ingredients such as pseudoephedrine, lithium batteries, and fertilizer, among others, a certain amount of the drug is produced locally in small, clandestine laboratories or through the use of a one-pot or “shake and bake” method where all ingredients are combined into one container (often a 2-liter or 20-ounce plastic soda bottle) and shaken (Blostein et al., 2009; Greene, Williams, & Wright, 2010). Clandestine labs create significant risks for persons who live in and around them due to the toxic fumes, chemical contamination, and risk of fires and explosions that are associated with this form of meth production, while the toxic residue from shake-and-bake production remaining in soda bottles is often dumped along roadways (Blostein et al., 2009; Greene, Williams, & Wright, 2010; Messina, Marinelli-Casey, West, & Rawson, 2007; Petit & Curtis, 1999). In 2018, the Indiana State Police (ISP) seized 192 clandestine meth labs and made 81 meth lab arrests. In the majority of the meth labs seized (n=164, 88%), the one-pot method was used. The number of meth labs seized in the state has seen a dramatic decline, particularly in the past two years with the number of labs seized in 2018 representing a 89% decrease from the peak number of seizures in 2013 (see Figure 6.10) (ISP, 2018).

Figure 6.9 Percentage of Treatment Episodes with Reported Prescription Stimulant Use and Dependence, Indiana and the United States (Treatment Episode Data Set, 2007-2016)



Source: SAMHSA, 2016

Figure 6.10 Number of Clandestine Methamphetamine Labs Seized and Number of Arrests Made at Methamphetamine Labs by the Indiana State Police (Indiana Meth Lab Statistics, 2009-2018)




Source: ISP, 2018

Children Taken from Methamphetamine Lab Homes

In addition to the health-related and criminal consequences, meth use can have serious social impacts on children and families in ways similar to other forms of substance abuse. These include contributing to increased interpersonal conflicts, violence, financial

problems, and poor parenting (Sommers, Baskin, & Baskin-Sommers, 2006). Other social effects of meth use include incarceration of parents and placement of children in protective custody. According to ISP data, the number of children who were taken from meth lab homes in Indiana peaked in 2013 (440 children), but dropped to 14 in 2018 (see Figure 6.11) (ISP, 2018).

Figure 6.11 Number of Indiana Children Taken by the Indiana State Police from Methamphetamine Lab Homes (Indiana Meth Lab Statistics, 2009-2018)



Source: ISP, 2018

APPENDIX 6A

Percentage of Indiana Students Reporting Monthly Cocaine and Methamphetamine Use, by Region and Grade (Indiana Youth Survey, 2018)

Cocaine									
	Indiana	North-west	North Central	North-east	West	Central	East	South-west	South-east
7th Grade	0.1	0.1	0.0	0.4	0.1	0.2	0.0	0.0	0.2
8th Grade	0.2	0.1	0.3	0.5	0.1	0.3	0.2	0.2	0.2
9th Grade	0.4	0.6	0.2	0.5	0.2	0.2	0.4	0.4	0.4
10th Grade	0.6	0.9	0.6	0.0	0.4	1.0	0.5	0.8	0.6
11th Grade	0.6	0.8	0.5	0.5	0.7	0.4	0.7	0.9	0.6
12th Grade	1.0	1.0	1.5	0.2	1.1	0.9	1.0	1.0	0.9
Methamphetamine									
	Indiana	North-west	North Central	North-east	West	Central	East	South-west	South-east
7th Grade	0.1	0.0	0.1	0.2	0.1	0.2	0.0	0.1	0.0
8th Grade	0.1	0.1	0.1	0.2	0.1	0.2	0.0	0.1	0.1
9th Grade	0.2	0.1	0.1	0.2	0.2	0.0	0.2	0.3	0.2
10th Grade	0.3	0.3	0.0	0.0	0.5	0.3	0.5	0.2	0.5
11th Grade	0.3	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.5
12th Grade	0.3	0.2	0.4	0.0	0.4	0.1	0.6	0.5	0.2

* Indicates a local rate that is significantly different from the overall state rate ($P < 0.05$).

Beginning in 2015, the Indiana Youth Survey stopped asking 6th grade students about cocaine and methamphetamine use.

Source: Gassman et al., 2018

APPENDIX 6B

Number of Treatment Episodes with Cocaine, Meth, and Prescription Stimulant Use and Dependence Reported at Treatment Admission in Indiana, by County (Substance Abuse Population by County/Treatment Episode Data Set, 2017)

County	Treatment Episodes	Cocaine Use		Cocaine Dependence		Meth Use		Meth Dependence		Rx Stimulant Use		Rx Stimulant Dependence	
	Total	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Adams	228	26	11.4%	<5	N/A	37	16.2%	17	7.5%	5	2.2%	<5	N/A
Allen	2050	543	26.5%	191	9.3%	205	10.0%	93	4.5%	35	1.7%	10	0.5%
Bartholomew	727	30	4.1%	<5	N/A	414	56.9%	249	34.3%	<5	N/A	<5	N/A
Benton	61	<5	N/A	<5	N/A	13	21.3%	5	8.2%	<5	N/A	<5	N/A
Blackford	144	16	11.1%	<5	N/A	68	47.2%	31	21.5%	8	5.6%	<5	N/A
Boone	167	15	9.0%	<5	N/A	51	30.5%	29	17.4%	5	3.0%	<5	N/A
Brown	72	6	8.3%	<5	N/A	36	50.0%	17	23.6%	<5	N/A	<5	N/A
Carroll	80	<5	N/A	<5	N/A	28	35.0%	13	16.3%	<5	N/A	<5	N/A
Cass	387	28	7.2%	7	1.8%	140	36.2%	92	23.8%	6	1.6%	<5	N/A
Clark	103	7	6.8%	<5	N/A	28	27.2%	14	13.6%	<5	N/A	<5	N/A
Clay	106	<5	N/A	<5	N/A	51	48.1%	40	37.7%	<5	N/A	<5	N/A
Clinton	147	7	4.8%	<5	N/A	62	42.2%	28	19.0%	<5	N/A	<5	N/A
Crawford	36	<5	N/A	<5	N/A	19	52.8%	12	33.3%	<5	N/A	<5	N/A
Daviess	202	<5	N/A	<5	N/A	98	48.5%	57	28.2%	5	2.5%	<5	N/A
Dearborn	497	61	12.3%	11	2.2%	76	15.3%	35	7.0%	11	2.2%	<5	N/A
Decatur	185	8	4.3%	<5	N/A	79	42.7%	44	23.8%	<5	N/A	<5	N/A
DeKalb	276	12	4.3%	<5	N/A	112	40.6%	71	25.7%	<5	N/A	<5	N/A
Delaware	843	101	12.0%	27	3.2%	235	27.9%	105	12.5%	17	2.0%	7	0.8%
Dubois	131	5	3.8%	<5	N/A	59	45.0%	25	19.1%	<5	N/A	<5	N/A
Elkhart	765	139	18.2%	63	8.2%	259	33.9%	160	20.9%	11	1.4%	7	0.9%
Fayette	322	12	3.7%	<5	N/A	130	40.4%	59	18.3%	7	2.2%	<5	N/A
Floyd	49	<5	N/A	<5	N/A	20	40.8%	10	20.4%	<5	N/A	<5	N/A
Fountain	59	5	8.5%	<5	N/A	27	45.8%	15	25.4%	<5	N/A	<5	N/A
Franklin	145	<5	N/A	<5	N/A	33	22.8%	12	8.3%	6	4.1%	<5	N/A
Fulton	188	8	4.3%	<5	N/A	73	38.8%	34	18.1%	<5	N/A	<5	N/A
Gibson	252	6	2.4%	<5	N/A	125	49.6%	74	29.4%	6	2.4%	<5	N/A
Grant	509	79	15.5%	20	3.9%	126	24.8%	40	7.9%	29	5.7%	9	1.8%
Greene	197	7	3.6%	<5	N/A	85	43.1%	44	22.3%	3	1.5%	<5	N/A
Hamilton	874	89	10.2%	28	3.2%	104	11.9%	49	5.6%	19	2.2%	<5	N/A
Hancock	398	43	10.8%	11	2.8%	73	18.3%	38	9.5%	6	1.5%	<5	N/A
Harrison	19	<5	N/A	<5	N/A	12	63.2%	6	31.6%	<5	N/A	<5	N/A
Hendricks	424	36	8.5%	14	3.3%	104	24.5%	57	13.4%	7	1.7%	<5	N/A
Henry	381	21	5.5%	5	1.3%	113	29.7%	65	17.1%	7	1.8%	<5	N/A
Howard	492	56	11.4%	15	3.0%	197	40.0%	65	13.2%	<5	N/A	<5	N/A
Huntington	136	8	5.9%	<5	N/A	38	27.9%	16	11.8%	<5	N/A	<5	N/A
Jackson	330	13	3.9%	<5	N/A	205	62.1%	146	44.2%	<5	N/A	<5	N/A
Jasper	93	8	8.6%	<5	N/A	28	30.1%	14	15.1%	<5	N/A	<5	N/A
Jay	122	15	12.3%	<5	N/A	41	33.6%	20	16.4%	<5	N/A	<5	N/A
Jefferson	307	13	4.2%	<5	N/A	153	49.8%	92	30.0%	<5	N/A	<5	N/A
Jennings	260	<5	N/A	<5	N/A	125	48.1%	87	33.5%	<5	N/A	<5	N/A
Johnson	424	37	8.7%	6	1.4%	137	32.3%	84	19.8%	8	1.9%	<5	N/A
Knox	352	5	1.4%	<5	N/A	160	45.5%	91	25.9%	13	3.7%	5	1.4%
Kosciusko	362	22	6.1%	6	1.7%	141	39.0%	67	18.5%	14	3.9%	<5	N/A

(continued on next page)

APPENDIX 6B (Continued from previous page)

County	Treatment Episodes	Cocaine Use		Cocaine Dependence		Meth Use		Meth Dependence		Rx Stimulant Use		Rx Stimulant Dependence	
	Total	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
LaGrange	115	5	4.3%	<5	N/A	62	53.9%	36	31.3%	<5	N/A	<5	N/A
Lake	1903	443	23.3%	180	9.5%	34	1.8%	9	0.5%	10	0.5%	<5	N/A
LaPorte	172	39	22.7%	17	9.9%	22	12.8%	7	4.1%	<5	N/A	<5	N/A
Lawrence	535	23	4.3%	5	0.9%	276	51.6%	166	31.0%	12	2.2%	<5	N/A
Madison	1205	140	11.6%	42	3.5%	372	30.9%	182	15.1%	27	2.2%	9	0.7%
Marion	4134	745	18.0%	309	7.5%	609	14.7%	280	6.8%	30	0.7%	7	0.2%
Marshall	137	9	6.6%	<5	N/A	35	25.5%	20	14.6%	<5	N/A	<5	N/A
Martin	60	<5	N/A	<5	N/A	26	43.3%	11	18.3%	<5	N/A	<5	N/A
Miami	249	8	3.2%	<5	N/A	89	35.7%	44	17.7%	5	2.0%	<5	N/A
Monroe	1428	124	8.7%	31	2.2%	557	39.0%	281	19.7%	26	1.8%	7	0.5%
Montgomery	304	23	7.6%	5	1.6%	113	37.2%	59	19.4%	14	4.6%	<5	N/A
Morgan	550	20	3.6%	<5	N/A	261	47.5%	172	31.3%	<5	N/A	<5	N/A
Newton	27	<5	N/A	<5	N/A	10	37.0%	<5	N/A	<5	N/A	<5	N/A
Noble	346	16	4.6%	7	2.0%	147	42.5%	105	30.3%	<5	N/A	<5	N/A
Ohio	33	<5	N/A	<5	N/A	9	27.3%	<5	N/A	<5	N/A	<5	N/A
Orange	135	<5	N/A	<5	N/A	60	44.4%	42	31.1%	<5	N/A	<5	N/A
Owen	187	8	4.3%	<5	N/A	81	43.3%	36	19.3%	<5	N/A	<5	N/A
Parke	65	<5	N/A	<5	N/A	23	35.4%	16	24.6%	<5	N/A	<5	N/A
Perry	94	<5	N/A	<5	N/A	49	52.1%	37	39.4%	<5	N/A	<5	N/A
Pike	46	<5	N/A	<5	N/A	18	39.1%	11	23.9%	<5	N/A	<5	N/A
Porter	202	35	17.3%	11	5.4%	10	5.0%	<5	N/A	<5	N/A	<5	N/A
Posey	156	<5	N/A	<5	N/A	75	48.1%	44	28.2%	6	3.8%	<5	N/A
Pulaski	82	6	7.3%	<5	N/A	21	25.6%	8	9.8%	5	6.1%	<5	N/A
Putnam	253	8	3.2%	<5	N/A	145	57.3%	67	26.5%	6	2.4%	<5	N/A
Randolph	167	25	15.0%	5	3.0%	66	39.5%	27	16.2%	<5	N/A	<5	N/A
Ripley	180	<5	N/A	<5	N/A	52	28.9%	24	13.3%	<5	N/A	<5	N/A
Rush	139	5	3.6%	<5	N/A	66	47.5%	46	33.1%	8	5.8%	<5	N/A
Saint Joseph	1351	379	28.1%	184	13.6%	196	14.5%	109	8.1%	20	1.5%	6	0.4%
Scott	212	9	4.2%	<5	N/A	129	60.8%	82	38.7%	<5	N/A	<5	N/A
Shelby	140	17	12.1%	<5	N/A	46	32.9%	29	20.7%	<5	N/A	<5	N/A
Spencer	132	<5	N/A	<5	N/A	73	55.3%	57	43.2%	<5	N/A	<5	N/A
Starke	77	5	6.5%	<5	N/A	38	49.4%	17	22.1%	<5	N/A	<5	N/A
Steuben	183	8	4.4%	<5	N/A	74	40.4%	42	23.0%	8	4.4%	<5	N/A
Sullivan	115	<5	N/A	<5	N/A	56	48.7%	33	28.7%	<5	N/A	<5	N/A
Switzerland	113	11	9.7%	<5	N/A	37	32.7%	21	18.6%	<5	N/A	<5	N/A
Tippecanoe	364	23	6.3%	7	1.9%	101	27.7%	51	14.0%	6	1.6%	<5	N/A
Tipton	46	<5	N/A	<5	N/A	12	26.1%	7	15.2%	<5	N/A	<5	N/A
Union	54	8	14.8%	<5	N/A	16	29.6%	9	16.7%	<5	N/A	<5	N/A
Vanderburgh	1218	63	5.2%	15	1.2%	575	47.2%	323	26.5%	29	2.4%	5	0.4%
Vermillion	132	<5	N/A	<5	N/A	75	56.8%	39	29.5%	<5	N/A	<5	N/A
Vigo	771	37	4.8%	11	1.4%	414	53.7%	256	33.2%	5	0.6%	<5	N/A
Wabash	299	7	2.3%	<5	N/A	93	31.1%	47	15.7%	5	1.7%	<5	N/A
Warren	26	<5	N/A	<5	N/A	7	26.9%	<5	N/A	<5	N/A	<5	N/A
Warrick	310	7	2.3%	<5	N/A	149	48.1%	94	30.3%	6	1.9%	<5	N/A
Washington	44	<5	N/A	<5	N/A	19	43.2%	10	22.7%	<5	N/A	<5	N/A

(continued on next page)

APPENDIX 6B (Continued from previous page)

County	Treatment Episodes	Cocaine Use		Cocaine Dependence		Meth Use		Meth Dependence		Rx Stimulant Use		Rx Stimulant Dependence	
	Total	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Wayne	558	102	18.3%	22	3.9%	114	20.4%	49	8.8%	6	1.1%	<5	N/A
Wells	156	18	11.5%	<5	N/A	38	24.4%	12	7.7%	6	3.8%	<5	N/A
White	125	7	5.6%	<5	N/A	49	39.2%	15	12.0%	<5	N/A	<5	N/A
Whitley	126	13	10.3%	<5	N/A	45	35.7%	22	17.5%	5	4.0%	<5	N/A
County Info Missing	41	6	14.6%	<5	N/A	12	29.3%	7	17.1%	<5	N/A	0	0.0%
Indiana	33,399	3,889	11.6%	1,255	3.8%	9,976	29.9%	5,383	16.1%	473	1.4%	72	0.2%

Notes: We defined dependence as “individuals in substance abuse treatment listing cocaine/meth/prescription stimulants as their primary substance at admission.”

We calculated the percentages by dividing the number of reported cocaine/meth/prescription stimulant use/dependence by the number of treatment episodes.

Information on treatment episodes <5 was suppressed due to confidentiality constraints.

Source: Indiana Family and Social Services Administration, 2018

REFERENCES, CHAPTER 6

- Blostein, P., Plaisier, B. R., Maltz, S. B., Davidson, S. B., Wideman, E. W., Feucht, E. C., & Vandenberg, S. L. (2009). Methamphetamine production is hazardous to your health. *The Journal of Trauma*, 66, 1712.
- Centers for Disease Control and Prevention. (1991-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (2007). *Methamphetamine use and risk for HIV/AIDS*. Retrieved from <http://www.cdc.gov/hiv/resources/factsheets/pdf/meth.pdf>
- Gassman, R., Jun, M., Samuel, S., Agle, J. D., Lee, L., & ... Wolf, J. (2018). *Indiana Youth Survey*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Greene, M. S., Williams, M. J., & Wright, E. R. (2010). *What's cooking? Meth use in Indiana*. Indianapolis, IN: Center for Health Policy, Department of Public Health, Indiana University School of Medicine.
- Halkitis, P. N., Parsons, J. T., & Stirratt, M. J. (2001). A double epidemic: Crystal methamphetamine
- Indiana Family and Social Services Administration. (2018). *Treatment Episode Data System (TEDS), 2018*. Indianapolis, IN: Indiana Family and Social Services Administration.
- Indiana State Police, Methamphetamine Suppression Section. (2018). *Indiana meth lab statistics, 2018*. Data received from First Sergeant Don McCay.
- Inter-university Consortium for Political and Social Research, University of Michigan. (2018). *Monitoring the Future (MTF)*. Retrieved from <http://www.monitoringthefuture.org/data/data.html>
- King, R. A., & Jun, M. K. (2018). *Indiana College Substance Use Survey, 2018*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey>
- Messina, N., Marinelli-Casey, P., West, K., & Rawson, R. (2007). Children exposed to methamphetamine use and manufacture. *Child Abuse & Neglect*, 38, 1872-1883.
- National Institute on Drug Abuse. (2016a). *Cocaine*. Retrieved from <https://d14rmgtrwzf5a.cloudfront.net/sites/default/files/1141-cocaine.pdf>
- National Institute on Drug Abuse. (2016b). *DrugFacts—Cocaine*. Retrieved from <http://www.drugabuse.gov/publications/drugfacts/cocaine>
- National Institute on Drug Abuse. (2017). *DrugFacts: Methamphetamine*. National Institutes of Health, U.S. Department of Health and Human Services. Retrieved from <http://www.drugabuse.gov/publications/drugfacts/methamphetamine>
- National Institute on Drug Abuse. (2018). *DrugFacts: Prescription stimulants*. Retrieved from <https://www.drugabuse.gov/publications/drugfacts/prescription-stimulants>
- Petit, M., & Curtis, P. (1999). *Child abuse and neglect: A look at the states*. Washington, DC: Children Welfare League of America.
- Sommers, I., Baskin, D., & Baskin-Sommers, A. (2006). Methamphetamine use among young adults: Health and social consequences. *Addictive Behaviors*, 31, 1469-1476.
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2016). *Treatment Episode Data Set -- Admissions (TEDS-A)*. Retrieved from <http://www.dasis.samhsa.gov/dasis2/teds.htm>
- Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>

7

MENTAL HEALTH AND SUICIDE IN INDIANA

INTRODUCTION

Good mental health is essential to a person's wellbeing. It affects our ability to adapt to change, cope with challenges, live productively, and have healthy relationships. Mental disorders are conditions characterized by alterations in thinking, mood, perception, and/or behavior (Office of Disease Prevention and Health Promotion, 2018). Mental illness collectively refers to all diagnosable mental disorders, including, but not limited to:

- Anxiety disorders (e.g., generalized anxiety disorder, phobias)
- Mood disorders (e.g., major depression, bipolar disorder)
- Psychotic disorders (e.g., schizophrenic spectrum and other psychotic disorders)
- Behavior disorders (e.g., ADHD, conduct disorder)
- Personality disorders (e.g., borderline or antisocial personality disorders)
- Substance-related and addictive disorders (e.g., alcohol and other substance use disorders) (Substance Abuse and Mental Health Services Administration, SAMHSA, 2019)

According to the Centers for Disease Control and Prevention (CDC, 2018b), more than 50% of Americans are diagnosed with a mental illness at some point during their lifetime, and 20% experience a mental disorder in a given year. Mental illness is associated with a number of other chronic diseases, as well as substance use (alcohol, tobacco, and drugs) and suicide (CDC, 2013; Kessler, 2004; SAMHSA, 2002, 2013).

The 2017 National Survey on Drug Use and Health (NSDUH) reported that of the 46.6 million U.S. adults who experienced a mental illness in the past year, 8.5 million (or 3.4%) also had a substance use disorder (SAMHSA, 2017a). Individuals diagnosed with co-occurring mental health and substance use disorders tend to have more complex problems, often resulting in a more chronic and persistent course of illness, poorer response to treatment, and higher rates of substance abuse relapse (Bradizza, Stasiewicz, & Paas, 2006; Davidson & White, 2007; Kessler, 2004).

For this chapter, we compiled available state-level data on indicators related to mental health. Definitions of specific terms used in this chapter can be found in Appendix 7A.


PREVALENCE OF PSYCHOLOGICAL DISTRESS IN INDIANA

National Survey on Drug Use and Health

The National Survey on Drug Use and Health (NSDUH) measures the prevalence of mental illness in the U.S. population. It defines 'any mental illness' (AMI) as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder; 'serious mental illness' (SMI) then refers to having a mental illness that results in serious functional impairment (2017a).

According to estimates from the 2017 NSDUH, one in five Indiana adults (20.9%) reported having any mental illness in the past year (95% CI [Confidence Interval]: 18.9-23.1), compared to 18.6% (95% CI: 18.2-18.9) of U.S. adults. Past-year prevalence rates for serious mental illness were similar in Indiana (5.2%, 95% CI: 4.4-6.2) and the nation (4.4%, 95% CI: 4.2-4.6). For AMI and SMI prevalence rates by age group, see Figure 7.1 (SAMHSA, 2017a).


Figure 7.1 Percentage of Indiana and U.S. Population (18 Years and Older) Reporting Any Mental Illness (AMI) or Serious Mental Illness (SMI) in the Past Year, by Age Group (National Survey on Drug Use and Health, 2017)



Source: SAMHSA, 2017a

Among adults ages 18 and older, past-year prevalence rates of AMI and SMI remained fairly stable between 2009 and 2017 (see Figure 7.2) (SAMHSA, 2017a).

Figure 7.2 Percentage of Indiana and U.S. Population (18 Years and Older) Reporting Any Mental Illness (AMI) or Serious Mental Illness (SMI) in the Past Year (National Survey on Drug Use and Health, 2009–2017)




Source: SAMHSA, 2017a

In 2017, 7.0% of Indiana adults (95% CI: 5.8-8.5) reported having had at least one major depressive episode (MDE) in the past year (U.S.: 6.1%, 95% CI: 5.8–6.3). For rates by age group, see Figure 7.3 (SAMHSA, 2017a).

The percentage of adults with a major depressive episode remained stable between 2008 and 2017 (see Figure 7.4) (SAMHSA, 2017a).


Figure 7.3 Percentage of Indiana and U.S. Population Reporting at Least One Major Depressive Episode in the Past Year, by Age Group (National Survey on Drug Use and Health, 2017)



Note: There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths ages 12 to 17 were not combined with data from persons ages 18 or older to produce the total MDE estimate.

Source: SAMHSA, 2017a

Figure 7.4 Percentage of Indiana and U.S. Population (18 Years and Older) Reporting at Least One Major Depressive Episode in the Past Year (National Survey on Drug Use and Health, 2008–2017)



Source: SAMHSA, 2017a

In 2017 more than 8.5 million U.S. adults (or 3.4%) had a co-occurring mental illness and substance use disorder; the prevalence rate was particularly high in young adults ages 18 to 25 (6.9%) (SAMHSA, 2017a). State-level estimates for co-occurring disorders are currently not available from the NSDUH.

Behavioral Risk Factor Surveillance System

According to the 2017 Behavioral Risk Factor Surveillance System (BRFSS), 23.5% of adults in Indiana reported ever being told that they had depression (U.S.: 20.5%). Among Hoosiers, having a history of depression was greatest among females, individuals who identified as multiracial, and individuals under the age of 65 (see Table 7.1) (CDC, 2017a).

Youth Risk Behavior Surveillance System

Based on the 2015 Youth Risk Behavior Surveillance System (YRBSS), the percentage of high school students who reported “stopping some of their normal activities during the past year due to feeling sad or hopeless almost every day for two weeks” did not differ significantly between Indiana and the nation (IN: 29.4%; U.S.: 29.9%). Rates were higher for females (39.2%) and students who self-identified as gay, lesbian, or bisexual (57.8%). For rates by student characteristics, see Table 7.2 (CDC, 1991-2015).

Table 7.1 Percentage of Indiana and U.S. Population (18 Years and Older) Reporting a History of Depression (Behavioral Risk Factor Surveillance System, 2017)

		Indiana (95% CI)
Gender	Male	17.0% (15.7-18.2)
	Female	29.6% (28.2-31.1)
Race/Ethnicity	White	24.9% (23.8-26.0)
	Black	18.8% (15.5-22.0)
	American Indian or Alaskan Native	20.3% (11.2-29.4)
	Multiracial	30.3% (22.7-37.9)
	Hispanic	11.1% (8.2-13.9)
Age Group	18-24	26.2% (22.4-30.0)
	25-34	25.9% (23.1-28.7)
	35-44	23.8% (21.2-26.3)
	45-54	24.8% (22.7-27.0)
	55-64	26.5% (24.6-28.5)
	65+	15.8% (14.6-17.0)
Total		23.5% (22.5-24.4)

Source: CDC, 2017a

Table 7.2 Percentage of Indiana and U.S. High School Students (Grades 9 through 12) Reporting Feeling Sad or Hopeless (Youth Risk Behavior Surveillance System, 2015)

		Indiana (95% CI)	U.S. (95% CI)
Gender	Male	19.8 (17.5–22.3)	20.3 (18.9–21.8)
	Female	39.2 (33.6–45.0)	39.8 (36.5–43.2)
Race/Ethnicity	White	28.4 (25.8–31.1)	28.6 (25.8–31.5)
	Black	31.2 (22.2–41.8)	25.2 (21.7–29.1)
	Hispanic	36.8 (27.8–46.8)	35.3 (32.3–38.4)
Grade	9th	26.9 (23.0–31.2)	28.4 (25.9–31.0)
	10th	33.3 (27.8–39.3)	29.8 (26.6–33.1)
	11th	31.8 (25.7–38.7)	31.4 (28.3–34.8)
	12th	26.0 (21.6–30.8)	30.0 (27.5–32.6)
Sexual Identity	Heterosexual	25.2 (22.5–28.0)	26.4 (24.6–28.4)
	Gay, Lesbian, or Bisexual	57.8 (44.8–69.8)	60.4 (55.1–65.4)
	Not Sure	44.6 (28.6–61.9)	46.5 (41.2–51.8)
Total		29.4 (27.0–31.9)	29.9 (27.0–31.9)

Source: CDC, 1991-2015

Physically and verbally threatening behaviors, most often in the form of bullying, have been linked to a number of mental health problems in youth, primarily depression and anxiety (CDC, 2018a). The YRBSS collects information on some of these indicators. According to 2015 findings:


- 6.6% of Indiana high school students (95% CI: 4.8–9.0) reported being threatened or injured on school property at least once with a weapon (U.S.: 6.0%, 95% CI: 5.2–6.8);
- 18.1% of Indiana high school students (95% CI: 15.0–21.6) reported being in a physical fight at least once (U.S.: 22.6%, 95% CI: 20.9–24.4);
- 15.7% of Indiana high school students (95% CI: 14.0–17.7) reported being electronically bullied (U.S.: 15.5%, 95% CI: 14.5–16.6); and

- 18.7% of Indiana high school students (95% CI: 16.1–21.5) reported being bullied on school property (U.S.: 20.2, 95% CI: 18.8–21.7) (CDC, 1991-2015).

Indiana Youth Survey

Results from the 2018 Indiana Youth Survey show that more than one-fifth of students in grades 6 through 12 reported feeling sad or hopeless. A substantial percentage of students also reported having considered suicide and even making a suicide plan in the past 12 months. For additional information, see Figure 7.5 (Gassman et al., 2018).

Figure 7.5 Percentage of Students who Experienced Feeling Sad or Hopeless, Considered Suicide, or Made a Suicide Plan in the Past 12 Months, Grades 6 through 12 (Indiana Youth Survey, 2018)



Source: Gassman et al., 2018

TREATMENT UTILIZATION

National Survey on Drug Use and Health

According to estimates from the 2017 NSDUH, 20.9% of adult Hoosiers experienced a mental illness in the past year (Figure 7.1), and 17.6% received mental health services (SAMHSA, 2017a).

Uniform Reporting System

In 2017, a total of 137,326 clients were served by the Indiana Division of Mental Health and Addiction (DMHA)—the state’s mental health authority. Of those, nearly all (136,699) were treated in community settings rather than state hospitals (1,066). The client population was predominately non-Hispanic (90.7%), white (76.9%), and slightly more than half were female (52.9%) (SAMHSA, 2017b).

Clients included children who met the federal definition for severe emotional disturbance (SED) and adults who met the federal definition for serious mental illness (SMI). Over one-fourth (26.0%) of adults served by DMHA received services for co-occurring mental illness and substance use disorders, as did 3.0% of the children (SAMHSA, 2017b). For more detailed client information, see Table 7.3.

SUICIDE

Suicide is a public health issue that is often associated with mental illness and substance use (CDC, 2017b; Lipari, Hughes, & Williams, 2016). Prior to actually making a suicide attempt, individuals may often spend significant amounts of time thinking about and planning how they might die by suicide.

Suicide is one of the top 10 leading causes of death for persons between the ages of 10 and 64 (National Center for Health Statistics, 2017). Although younger individuals are more likely to think about suicide, suicide deaths most frequently occur in adults between the ages of 45 and 54 (CDC, 2017b).

National Survey on Drug Use and Health

According to 2017 NSDUH findings, 5.0% of Indiana adults (95% CI: 4.2–6.1) reported having serious thoughts of suicide in the past year; an estimate similar to the U.S. rate of 4.2% (95% CI: 4.0–4.4). This was particularly prevalent among young adults ages 18 to

Table 7.3 Demographic Characteristics of Adults with SMI and Children with SED Served by the Indiana Division of Mental Health and Addiction, FY 2017

Gender	Male	47.1%
	Female	52.9%
Race	White	76.9%
	Black	14.6%
	Other/Unknown	8.5%
Ethnicity	Hispanic	6.3%
Age Group	Children 0-17	39.7%
	Adults 18+	60.3%
Medicaid Status	Medicaid only	69.0%
	Both Medicaid and other funds	19.0%
	Non-Medicaid	12.0%
Total		137,326 (100%)

Source: SAMHSA, 2017b

25 (IN: 11.1%, 95% CI: 9.0–13.5; U.S.: 9.6%, 95% CI: 9.2–10.1) (SAMHSA, 2017a).

Youth Risk Behavior Surveillance System

Based on estimates from the 2015 YRBSS, nearly one in ten high school students attempted suicide in the past year. The overall percentages were similar in Indiana (9.9%) and the U.S. (8.6%). Rates were particularly high for students who self-identified as gay, lesbian, or bisexual (34.2%). For prevalence rates by gender, race/ethnicity, sexual identity, and grade level, see Table 7.4 (CDC, 1991-2015).

Suicide Mortality


Suicide deaths both nationally and in Indiana have increased significantly since 1999 (IN: 10.4; U.S.: 10.5, per 100,000 population). According to 2017 estimates, Indiana’s age-adjusted suicide mortality rate of 16.4 per 100,000 population (95% CI: 15.4–17.4) was significantly higher than the U.S. rate of 14.5 (95% CI: 14.3–14.6). For 10-year trends, see Figure 7.6. Most suicide deaths occurred in males, whites, and non-Hispanics (see Table 7.5). For county-level age-adjusted annual suicide mortality rates, refer to Map 7.1 (CDC, 1999-2017).

Table 7.4 Percentage of Indiana and U.S. High School Students (Grades 9 through 12) Reporting Attempting Suicide in the Past Year (Youth Risk Behavior Surveillance System, 2015)

		Indiana (95% CI)	U.S. (95% CI)
Gender	Male	8.7 (6.0–12.5)	5.5 (4.7–6.4)
	Female	10.9 (8.3–14.1)	11.6 (9.7–13.7)
Race/Ethnicity	White	8.7 (6.5–11.5)	6.8 (5.5–8.4)
	Black	14.5 (8.8–23.1)	8.9 (6.7–11.9)
	Hispanic	15.5 (8.9–25.8)	11.3 (9.9–13.0)
Grade	9th	12.8 (7.7–12.7)	9.9 (8.5–11.5)
	10th	11.4 (8.6–14.9)	9.4 (7.6–11.6)
	11th	10.0 (6.4–15.2)	8.0 (6.8–9.5)
	12th	5.0 (2.7–9.0)	6.2 (4.9–7.9)
Sexual Identity	Heterosexual	6.8 (5.0–9.2)	6.4 (5.6–7.3)
	Gay, Lesbian, or Bisexual	34.2 (27.5–41.5)	29.4 (25.7–33.3)
	Not Sure	17.6 (7.5–35.9)	13.7 (10.0–18.5)
Total		9.9 (7.7–12.7)	8.6 (7.6–9.6)

Source: CDC, 1991-2015

Figure 7.6 Age-Adjusted Suicide Mortality Rate per 100,000 Population in Indiana and the United States (CDC WONDER, 2008–2017)



Source: CDC, 2008-2017

Table 7.5 Age-Adjusted Suicide Mortality Rate per 100,000 Population in Indiana and the United States (CDC WONDER, combined data from 1999-2017)

		Indiana (95% CI)	U.S. (95% CI)
Gender	Male	27.3 (25.4–29.1)	22.4 (22.1–22.6)
	Female	6.0 (5.1–6.8)	6.1 (6.0–6.2)
Race	White	17.5 (16.4–18.6)	15.8 (15.7–16.0)
	Black	8.8 (6.7–11.3)	6.6 (6.4–6.9)
	Asian or Pacific Islander	Unreliable (2.6–13.3)	6.6 (6.3–7.0)
	American Indian or Alaska Native	N/A	13.5 (12.4–14.6)
Ethnicity	Hispanic	7.9 (4.7–12.3)	6.9 (6.7–7.1)
	Not Hispanic	17.0 (15.9–18.0)	15.3 (15.2–15.5)
Total		16.3 (15.3–17.3)	14.0 (13.9–14.1)

Source: CDC, 1999-2017

Appendix 7A Definitions and Explanations

Any Mental Illness (AMI): “AMI among adults aged 18 or older is defined as currently or at any time in the past 12 months having had a diagnosable mental, behavioral, or emotional disorder (excluding developmental and substance use disorders) of sufficient duration to meet diagnostic criteria specified within the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)” (SAMHSA, 2017a).

Serious Mental Illness (SMI): “SAMHSA defined SMI as persons aged 18 or older who currently or at any time in the past year have had a diagnosable mental, behavioral, or emotional disorder (excluding developmental and substance use disorders) of sufficient duration to meet the criteria specified within DSM-IV that has resulted in serious functional impairment, which substantially interferes with or limits one or more major life activities” (SAMHSA, 2017a).

Major Depressive Episode (MDE): “MDE, as defined in NSDUH, is based on the definition of MDE in the DSM-IV (APA, 1994) and is measured for the lifetime and past year periods. Lifetime MDE is defined as having at least five or more of nine symptoms of depression in the same 2-week period in a person’s lifetime, in which at least one of the symptoms was a depressed mood or loss of interest or pleasure in daily activities. Respondents who had MDE in their lifetime were defined as having past

year MDE if they had a period of depression lasting 2 weeks or longer in the past 12 months while also having some of the other symptoms of MDE. It should be noted that, unlike the DSM-IV criteria for MDE, no exclusions were made in NSDUH for depressive symptoms caused by medical illness, bereavement, or substance use disorders” (SAMHSA, 2017a).

Depression: “Has a doctor, nurse, or other health professional EVER told you that you had...a depressive disorder, including depression, major depression, dysthymia, or minor depression?” (CDC, 2017a).

Feeling Sad or Hopeless:

- a) “Felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)” (CDC, 1991-2015).
- b) “During the past 12 months, did you ever feel so sad or hopeless almost every day for 2 weeks or more in a row that you stopped doing some usual activities?” (Gassman et al., 2018).

Suicide Attempts: “Attempted suicide one or more times during the 12 months before the survey” (CDC, 2017b).

Suicide Deaths: Suicide (intentional self-harm) deaths include ICD-10 codes U03.0 (Terrorism involving explosions and fragments), U03.9 (Terrorism by other and unspecified means), X60-X84 (Intentional self-harm).

REFERENCES, CHAPTER 7

- Bradizza, C. M., Stasiewicz, P. R., & Paas, N. D. (2006). Relapse to alcohol and drug use among individuals diagnosed with co-occurring mental health and substance use disorders: A review. *Clinical Psychology Review, 26*(2), 162–178. doi:10.1016/j.cpr.2005.11.005.
- Centers for Disease Control and Prevention. (1991-2015). *Youth Risk Behavior Surveillance System (YRBSS)*. Retrieved from <http://nccd.cdc.gov/youthonline>
- Centers for Disease Control and Prevention. (1999-2017). *CDC WONDER underlying causes of death (detailed mortality)*. Retrieved from <http://wonder.cdc.gov/>
- Centers for Disease Control and Prevention. (2013). Vital signs: Current cigarette smoking among adults aged ≥18 years with mental illness – United States, 2009-2011. *MMWR* 2013; 62(05): 81-87
- Centers for Disease Control and Prevention. (2017a). *Behavioral Risk Factor Surveillance System (BRFSS) prevalence & trends data*. Retrieved from <http://www.cdc.gov/brfss/brfssprevalence/index.html>
- Centers for Disease Control and Prevention. (2017b). *Suicide Prevention*. Retrieved from <http://www.cdc.gov/ViolencePrevention/suicide/index.html>
- Centers for Disease Control and Prevention. (2018a). *Bullying Research*. Retrieved from <http://www.cdc.gov/violenceprevention/youthviolence/bullyingresearch/index.html>
- Centers for Disease Control and Prevention. (2018b). *Learn About Mental Health*. Retrieved from <https://www.cdc.gov/mentalhealth/learn/index.htm>
- Davidson, L., & White, W. (2007). The Concept of Recovery as an Organizing Principle for Integrating Mental Health and Addiction Services. *The Journal of Behavioral Health Services and Research, 34*(2), 109–120. doi:10.1007/s11414-007-9053-7.
- Gassman, R., Jun, M., Samuel, S., Agle, J. D., Lee, J., & Wolf, J. (2018). *Indiana Youth Survey*. Indiana Prevention Resource Center, Indiana University. Retrieved from <http://inys.indiana.edu/survey-results>
- Kessler, R. C. (2004). The epidemiology of dual diagnosis. *Biological Psychiatry, 56*(10), 730–737. doi:10.1016/j.biopsych.2004.06.034.
- Lipari, R. N., Hughes, A., & Williams, M. (2016). *State estimates of past year serious thoughts of suicide among young adults: 2013-2014*. The CBHSQ Report: - June 16, 2016. Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Rockville, MD.
- National Center for Health Statistics (2017). *10 Leading Causes of Death--2015*. National Vital Statistics System, Office of Statistics and Programming, National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.
- Office of Disease Prevention and Health Promotion. (2018). *Healthy People 2020: Mental Health and Mental Disorders*. Available at <https://www.healthypeople.gov/2020/topics-objectives/topic/mental-health-and-mental-disorders>
- Substance Abuse and Mental Health Services Administration. (2002). *Report to Congress on the prevention and treatment of co-occurring substance abuse disorders and mental disorders*. Washington DC: United States Department of Health and Human Services.
- Substance Abuse and Mental Health Services Administration. (2019). *Mental Disorders*. Available at <https://www.samhsa.gov/disorders/mental>

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2013). The NSDUH Report Data Spotlight: *Adults with Mental Illness or Substance Use Disorder Account for 40 Percent of All Cigarettes Smoked*. Rockville, MD.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017a). *National Survey on Drug Use and Health (NSDUH)*. Retrieved from <https://www.samhsa.gov/data/population-data-nsduh>

Substance Abuse and Mental Health Services Administration, Center for Mental Health Services (SAMHSA/CMHS) (2017b). *Indiana 2017 Mental Health National Outcome Measures (NOMS): SAMHSA Uniform Reporting System*. Retrieved from <https://www.dasis.samhsa.gov/dasis2/urs.htm>

8

METHODS

This annual report describes the consumption and consequences of alcohol, tobacco, and other drugs in Indiana. We analyzed patterns within Indiana's general population, and compared them to patterns found among the U.S. population. Based on discussions with the State Epidemiological Outcomes Workgroup (SEOW), we have reviewed consumption and consequences data for the following drugs: alcohol, tobacco, marijuana, opioids, and stimulants. Additionally, we examined indicators of mental health and suicide in Indiana.

Our research team completed statistical analyses on publicly available local and national data sets using Statistical Package for the Social Sciences (SPSS) and Statistical Analysis System (SAS) software. For surveys that do not have publicly available data sets, we conducted statistical analyses using online analysis software and/or analysis tables provided by the agencies that conducted the data collection. Whenever possible, we made statistical comparisons across gender, racial/ethnic, and age groups for both drug-consumption behaviors and drug-use consequences. For all comparisons, a *P* value of .05 or less, or the 95 percent Confidence Interval (CI) was used to determine statistical significance.¹

Prevalence rates and other statistics may be presented somewhat differently across chapters, depending on the data sources that provided the information.

We used two guidelines to determine potential priorities. The first guideline was statistical significance. Statistical significance is a mathematical concept used to determine whether differences between groups are true or due to chance. Significance in this context does not necessarily mean "meaningful" and does not convey practical or clinical importance. Specific drug consumption and consequence patterns that place Indiana statistically significantly higher than the United States were used as markers for areas that could potentially benefit from intervention.

The second guideline was clinical or substantive significance; i.e., consumption behaviors or drug-use

consequences that are trending toward a higher frequency within a particular group of Hoosiers, such as a specific gender, race/ethnicity, or age group.

DATA SOURCES

The data for these analyses were gathered from various publicly available federal, state, and local-level surveys and administrative data sets. In order to compare Indiana with the nation as a whole and to determine trends in drug use and drug-related consequences over time, we selected, whenever possible, surveys and data sources that had at least two years' worth of data available. In all cases, the most recent findings were included.

CONSIDERATIONS

This report relies primarily on the data sources listed below. These are either 1) publicly available sources that our researchers could access and analyze for this year's state epidemiological report or 2) agency data sources that were provided specifically to the SEOW. Because of the nature of the available data, there are significant limitations to the interpretations presented:

- Consistent comparisons across data sources are not always possible due to the nature of the survey questions asked and information gathered.
- Inconsistencies may occur within classifications of demographic characteristics (e.g., age ranges, racial categories, grade levels).
- Timeframes may be inconsistent for comparisons across substances and data sources (e.g., some data have longer gaps than others before they are made publicly available).
- State-level prevalence rates presented in national surveys are often estimated using statistical algorithms.
- Due to the reporting requirements for national databases, the data may not be representative of the actual population of either the state or the nation..

¹Throughout the chapters, we use the terms "significant," "significantly different," or "statistically different" to report a statistically significant difference between groups.

In future editions of this report, we will expand the data analysis as additional data sources are made available to the SEOW data analysis team.

SEOW DATA SOURCES LIST

Following is a list of the data sources used in this report.

Alcohol-Related Disease Impact (ARDI) Database

The Centers for Disease Control and Prevention's (CDC) ARDI software generates estimates of alcohol-related deaths and years of potential life lost (YPLL) due to alcohol consumption. To do this, ARDI either calculates estimates or uses predetermined estimates of alcohol-attributable fractions (AAFs)—that is, the proportion of deaths from various causes that are due to alcohol. These AAFs are then multiplied by the number of deaths caused by a specific condition (e.g., liver cancer) to obtain the number of alcohol-attributable deaths. Reports can be generated based on national or state-level data.

Description: ARDI provides state and national estimates on alcohol-related deaths and years of potential life lost (YPLL) based on alcohol-attributable fractions.

Sponsoring Organization/Source: CDC.

Geographic Level: National and state levels.

Availability: The database can be accessed at http://nccd.cdc.gov/DPH_ARDI/default/default.aspx.

Trend: Pooled data averages from 2006–2010.

Strengths/Weaknesses: ARDI may underestimate the actual number of alcohol-related deaths and years of potential life lost.

Automated Reporting Information Exchange System (ARIES)

The Indiana State Police's ARIES is a central repository for all vehicle collisions reported in the state of Indiana, with and without alcohol involvement. Information on fatal accidents contained in the system is submitted to the national Fatality Analysis Reporting System.

Description: ARIES contains data on vehicle crashes with and without alcohol involvement.

Sponsoring Organization/Source: Indiana State Police (ISP).

Geographic Level: State and county levels.

Availability: Upon request from the ISP.

Trend: Annual; most recent data from 2017.

Strengths/Weaknesses: The data are in aggregate format; comparisons by demographic variables such as age, gender, and race/ethnicity are not possible.

Behavioral Risk Factor Surveillance System (BRFSS)

The CDC conducts the BRFSS annually with the assistance of health departments in all 50 states and the District of Columbia, Puerto Rico, Guam, and the U.S. Virgin Islands. BRFSS asks respondents ages 18 and older questions about health-related behaviors, including alcohol consumption and tobacco use. BRFSS results are available at the national and state levels as well as for selected metropolitan/micropolitan areas. BRFSS data allow for statistical comparisons across gender, age, race/ethnicity, educational attainment, and income level.

The BRFSS has traditionally used random-digit-dial telephone sampling of households with landline telephones. However, the increasing percentage of households abandoning their landline telephones for cell phones has significantly eroded the population coverage provided by landline-based surveys to 70% of the U.S. household population. To meet challenges for increasing non-coverage and decreasing response rates due to cell-phone-only households, BRFSS has expanded its traditional methodology to a dual frame survey of landline and cell phone numbers and has introduced a new weighting method called iterative proportional fitting, or raking. It would not be appropriate to directly compare estimates prior to 2011 with later estimates, due to different data adjustment methods and different sampling frames.

Description: BRFSS is an annual state health survey that monitors risk behaviors, including alcohol and tobacco consumption, related to chronic diseases, injuries, and death.

Sponsoring Organization/Source: CDC.

Geographic Level: National and state levels; selected metropolitan/micropolitan areas.

Availability: National and state data are available from the CDC at <https://www.cdc.gov/brfss/brfssprevalence>.

Trend: Annual; most recent data from 2017.

Strengths/Weaknesses: CDC consistently works to test and improve BRFSS methodology in an effort to make findings result in more valid and reliable data for public health surveillance. Due to substantial changes in methodology starting with the 2011 survey, comparison

of current estimates with estimates from previous years would not be appropriate.

Hospital Discharge Data

The Indiana State Department of Health (ISDH) collects information on inpatients discharged from hospitals in Indiana. The data are publicly available in aggregate format and include information on hospitals, principal diagnoses and procedures, length of stay, total charges, etc.

Description: Hospital discharge data are publicly available in aggregate format. Dataset can be queried by primary diagnosis (ICD-10-CM codes), e.g., for alcohol- and drug-induced diseases.

Sponsoring Organization/Source: ISDH.

Geographic Level: Indiana.

Availability: Annual data are available at <http://www.in.gov/isdh/20624.htm>.

Trend: Annual; most recent data from 2017.

Strengths/Weaknesses: The data are in aggregate format; comparisons by demographic variables such as age, gender, and race/ethnicity are not possible. Comparisons to years prior to 2016 are not possible due to the ICD-9-CM to ICD-10-CM switch that occurred on October 1, 2015.

Indiana Adult Tobacco Survey (IATS)

The Indiana Adult Tobacco Survey (IATS), a survey by the ISDH Tobacco Prevention and Cessation Commission (TPCC), collects information on tobacco use, cessation attempts, and other related issues among Hoosiers ages 18 and older. The survey uses a random-sampling design; African-American and Hispanic adults as well as residents in more rural regions of the state are oversampled. Data are available by gender, race/ethnicity, age group, income level, educational attainment, Indiana region, health insurance type, and number of children in household.

Description: This survey measures tobacco use among Indiana adults, and includes items on tobacco use, cessation, secondhand smoke, and awareness.

Sponsoring Organization/Source: ISDH/TPCC.

Geographic Level: Indiana.

Availability: Datasets can be requested from ISDH/TPCC; reports are available at <http://www.in.gov/isdh/tpc/2343.htm>.

Trend: Biannual; most recent data from 2017.

Strengths/Weaknesses: IATS uses a random-sample design, making findings representative of all Hoosier

adults. Oversampling of African-American and Hispanic adults, as well as residents in more rural regions, provides more robust estimates for these population groups.

Indiana College Substance Use Survey

The Indiana College Substance Use Survey was developed in 2009 by the Indiana Collegiate Action Network (ICAN) and the Indiana Prevention Resource Center (IPRC), with input from Indiana institutions of higher education and the Indiana State Epidemiological Outcomes Workgroup. The instrument was designed to assess prevalence of alcohol, tobacco, and other drug use; consequences of use; alcohol availability; and student perceptions of peer behaviors among Indiana college students. Information is available by gender, age category (under 21 vs. 21 or over), and type of institution (private vs. public). All two- and four-year colleges in Indiana are invited to participate in the survey. Results are based on nonrandom sampling and are not representative of all college students in Indiana.

Description: The survey measures the prevalence of alcohol, tobacco, and other drug use; consequences of use; alcohol availability; and student perceptions of peer behaviors among Indiana college students.

Sponsoring Organization/Source: Indiana Collegiate Action Network (ICAN); Indiana Prevention Resource Center (IPRC).

Geographic Level: Indiana.

Availability: Annual data are available at <http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey>.

Trend: Annual; most recent data from 2018.

Strengths/Weaknesses: The survey utilizes a nonrandom sampling design; results, therefore, are not representative of all college students in Indiana.

Indiana Meth Lab Statistics

The Indiana State Police (ISP) collects data on clandestine meth lab seizures in the state, including number of meth labs seized, number of arrests made during lab seizures, and the number of children located at/rescued from meth labs. The information is then submitted to National Clandestine Laboratory Seizure System, a database maintained by the U.S. Drug Enforcement Administration and the El Paso Intelligence Center. State and county-level information can be requested from the ISP.

Description: ISP collects meth lab incidence data

including: Number of meth labs seized, number of arrests made during lab seizures, and the number of children located at/rescued from meth labs.

Sponsoring Organization/Source: ISP.

Geographic Level: State and county level.

Availability: Indiana data from ISP are available on request; national data can be accessed at <http://www.dea.gov/resource-center/meth-lab-maps.shtml>.

Trend: Annual; most recent data from 2018.

Strengths/Weaknesses: The data include all meth incidents, including labs, “dumpsites,” or “chemical and glassware” seizures.

Indiana Mortality Data and National Vital Statistics System (NVSS)

NVSS is a CDC-maintained data system that provides information on mortality rates by cause of death as coded in the World Health Organization’s International Classification of Diseases, 10th Edition (ICD-10). Health departments in the 50 states, the District of Columbia, and U.S. territories provide CDC with data on deaths throughout the country. Using the query system on CDC’s website (CDC WONDER), researchers can compute mortality rates for deaths due to diseases and events associated with alcohol, tobacco, and other drug use (e.g., cirrhosis, lung cancer, heart disease, suicide, homicide, etc.) at the national, state, and county level. The system also allows for comparisons across gender, age, and racial groups. Indiana mortality data can also be requested directly from the Indiana State Department of Health (ISDH).

Description: NVSS contains mortality data from all U.S. states; the online database can be queried on number of deaths and death rates from alcohol- and drug-related causes. Indiana data can also be requested directly from ISDH.

Sponsoring Organization/Source: CDC’s National Center for Health Statistics; ISDH.

Geographic Level: National, state, and county levels.

Availability: National mortality data can be accessed by underlying cause of death (ICD-10 codes) from CDC at <http://wonder.cdc.gov/mortSQL.html>; state data are available on request from ISDH.

Trend: Annual; most recent data from 2017.

Strengths/Weaknesses: The strengths of the NVSS include availability of multiple years of data and the relatively large number of American Indian, Alaska Native, and other Native American respondents. However, a

primary weakness of the data is the quality of the race/ethnicity information, particularly for the American Indian/Alaska Native category, as data quality checks of the racial/ethnic distribution of the deceased in this category are lower than the distribution represented in Census estimates.

Indiana Scheduled Prescription Electronic Collection & Tracking (INSPECT)

INSPECT is the state’s prescription drug monitoring program. The secure database collects basic demographic information on the patient, the type of controlled substance prescribed, the prescribing practitioner, and the dispensing pharmacy. Each time a controlled substance is dispensed, the dispenser (e.g., pharmacy, physician, etc.) is required to submit the information to INSPECT. The program was designed to help address problems of prescription drug abuse and diversion in Indiana. By compiling controlled substance information into an online database, INSPECT performs two critical functions: (1) maintaining a warehouse of patient information to assist healthcare professionals in making treatment decisions; and (2) providing an important investigative tool for law enforcement to help prevent the possible diversion of controlled substances.

Description: INSPECT is Indiana’s prescription drug monitoring program; the online database collects information each time a controlled substance is dispensed.

Sponsoring Organization/Source: Indiana Professional Licensing Agency (IPLA).

Geographic Level: Indiana and counties.

Availability: Number and rate of opioid dispensations aggregated at the county and Indiana-level is available from ISDH at https://gis.in.gov/apps/isdh/meta/stats_layers.htm.

Trend: Annual; most recent 2018.

Strengths/Weaknesses: Data collection is statewide, and licensed dispensers (e.g., pharmacies, physicians) are required to submit information each time a controlled substance is dispensed. Dispensations aggregated at the county-level are approximate as some dispensations do not have a designated county FIPS code.

Indiana Youth Survey (INYS)

The Indiana Youth Survey is an annual school-based assessment conducted by the Indiana Prevention Resource Center (IPRC) and funded through the Indiana

Family and Social Services Administration/Division of Mental Health and Addiction. The survey is designed to monitor patterns of alcohol, tobacco, and other drug use; gambling behaviors; and risk and protective factors among Indiana middle and high school students, grades 6 through 12.

Caution is needed when comparing findings to previous years due to changes made to the survey in 2015. These changes, in addition to a revised cleaning methodology, make it difficult to draw accurate comparisons to the prevalence data from previous years.

The Indiana Youth Survey uses a convenience sampling design; i.e., the survey is open to all Indiana schools or school corporations, resulting in a large number of usable responses. However, the rate of participation varies widely across regions. In 2016, INYS also incorporated a random sampling process. The advantage of simultaneously collecting both random and convenience samples is that state-level estimates can be interpreted with greater confidence, even in areas with low participation rates.

INYS results are often compared to findings from the Monitoring the Future (MTF) survey conducted by the National Institute on Drug Abuse (<http://www.monitoringthefuture.org/data/data.html>). MTF is an ongoing study of youth behaviors, attitudes, and values about substance use; students in 8th, 10th, and 12th grades are surveyed annually.

Description: The IPRC manages the Indiana Youth Survey. The survey is administered to students (6th through 12th graders) annually in a number of schools throughout the state.

Sponsoring Organization/Source: IPRC; Indiana Family and Social Services Administration (FSSA)/Indiana Division of Mental Health and Addiction (DMHA).

Geographic Level: Indiana state and regions.

Availability: Reports with data tables are available at <http://inys.indiana.edu/survey-results>.

Trend: Annual; most recent data from 2018.

Strengths/Weaknesses: School-specific survey results are valuable to participating schools and provide statewide prevalence estimates. Due to changes made to the survey, data cannot be compared to findings from previous years (prior to 2015).

Indiana Youth Tobacco Survey (IYTS)

The CDC developed the National Youth Tobacco Survey as a way to estimate the current use of tobacco products

among middle school and high school students in the United States. Student respondents are asked to describe their lifetime, annual, and current use of cigarettes and other tobacco products. The Indiana State Department of Health's Tobacco Prevention and Cessation Commission (ISDH/TPCC) oversees Indiana's version of the survey, which includes CDC core and recommended questions, as well as state-specific items. IYTS is conducted every other year (even years); findings allow comparisons across gender, race/ethnicity, and grade levels.

Description: IYTS is Indiana's adapted version of CDC's NYTS. The surveys collect data from students in grades 6 through 12 on all types of tobacco use, exposure to secondhand smoke, and access to tobacco.

Sponsoring Organization/Source: CDC; ISDH/TPCC.

Geographic Level: Indiana.

Availability: Data are available on request from TPCC, and annual reports can be accessed at <http://www.in.gov/isdh/tpc/2343.htm>. National data are available at http://www.cdc.gov/tobacco/data_statistics/surveys/NYTS/.

Trend: Biannual; most recent data from 2016.

Strengths/Weaknesses: The IYTS provides detailed statewide information regarding youth knowledge, attitudes, and behaviors. However, county-level data are not available.

National Survey on Drug Use and Health (NSDUH)

NNSDUH is a national survey funded by the Substance Abuse and Mental Health Services Administration (SAMHSA) and designed to monitor patterns and track changes in substance use among U.S. residents 12 years of age and older. The survey asks respondents to report on use and misuse of substances including alcohol, tobacco, marijuana, cocaine, heroin, and prescription medications. Additionally, NSDUH asks respondents whether they received treatment for drug abuse or drug dependence during the past (prior) year. The survey also includes several modules of questions that focus on mental health issues.

Prevalence rates for substance use and specific mental health indicators are provided for the nation and each state. Raw data files from NSDUH surveys are publicly available; however, they do not allow for comparisons among states because NSDUH eliminates state identifiers in the process of preparing public-use data files. Tables with prevalence numbers and rates are prepared by SAMHSA's Center for Behavioral Health

Statistics and Quality and can be accessed online. Data reports are available since 1994. There is usually a two-year delay from the time of data collection to its availability.

In 2015, several changes were made to the NSDUH questionnaire and data collection process, causing some estimates not to be comparable with estimates from previous years. Items affected by these changes included binge drinking and prescription drug misuse. Due to these revisions, 2015 and later estimates cannot be compared to earlier years.

Description: NSDUH provides national and state-level estimates on the use of alcohol, tobacco, and illicit drugs (including nonmedical prescription drug use), as well as mental health indicators in the general population ages 12 and older.

Sponsoring Organization/Source: SAMHSA.

Geographic Level: National and state; some sub-state data are available using small-area estimation techniques.

Availability: National and state data tables are available at the NSDUH website at <http://www.samhsa.gov/data/population-data-nsduh>.

Trend: Annual; most recent data from 2017.

Strengths/Weaknesses: State-level data do not allow for comparisons by gender or race/ethnicity.

Treatment Episode Data Set (TEDS)

TEDS is a national database maintained by Substance Abuse and Mental Health Services Administration (SAMHSA) that records information about individuals entering treatment for substance abuse and/or dependence. State mental health departments submit data to TEDS on an annual basis. The information reported in TEDS includes age, race, ethnicity, gender, and other demographic characteristics, as well as information on the use of various substances. The data represent admissions rather than individuals, thus individuals may be admitted to treatment more than once in a given year. TEDS data become publicly available approximately two years after the information is gathered. The format of the TEDS data allows for comparisons between Indiana and the United States by gender, race, and age group.

County-level TEDS data for Indiana are available from the Indiana Family and Social Services Administration (FSSA), Division of Mental Health and Addiction (DMHA). While TEDS data can provide some information on drug use and abuse patterns both nationally and at the state level, the population on which the data are based

may not be representative of all individuals in drug and alcohol treatment. For Indiana, TEDS data are limited to information on individuals entering substance abuse treatment who are 200% below the federal poverty level and receive state-funded treatment.

Description: TEDS provides information on demographic and substance abuse characteristics of individuals in alcohol and drug abuse treatment. Data are collected by treatment episode. A treatment episode is defined as the period from the beginning of treatment services (admission) to termination of services.

Sponsoring Organization/Source: SAMHSA; FSSA/DMHA.

Geographic Level: National, state, and county-level.

Availability: National and state TEDS data were acquired from SAMHSA's Drug & Alcohol Services Information System at <http://www.dasis.samhsa.gov/dasis2/teds.htm>; county-level data available from FSSA upon request.

Trend: Annual; most recent data from 2016 (from SAMHSA) and 2018 (from DMHA).

Strengths/Weaknesses: In Indiana, these data are not representative of the state as a whole, as only individuals who are at or below the 200% poverty level are eligible for treatment at state-registered facilities.

Youth Risk Behavior Surveillance System (YRBSS)

The YRBSS is a national survey of health-related behaviors among students in grades 9 through 12. The CDC conducts the survey biannually with the cooperation of state health departments throughout the nation. Student respondents are asked to describe whether they have engaged in numerous behaviors that could pose a danger to their health, including the use of alcohol, tobacco, and other drugs. CDC's online database allows comparisons between Indiana and the United States on gender, race/ethnicity, and grade level. Data for the YRBSS are available every other year (odd years), with a one-year lag between the end of data collection and the publication of results. Though YRBSS data for some states are available from 1991, Indiana started participating in data collection in 2003. Availability of state-level results is dependent upon sufficient participation to achieve an adequate response rate to weight the data.

Description: This biannual national survey monitors health risks and behaviors among youth in grades 9 through 12.

Sponsoring Organization/Source: CDC.

Geographic Level: National and state level.

Availability: National and state-level data are downloadable from selected published tables on the CDC website at <http://nccd.cdc.gov/YouthOnline/App/Default.aspx>.

Trend: Biannual; most recent data from 2015.

Strengths/Weaknesses: Availability of state-level results is dependent upon sufficient participation; Indiana's response rates in 2013 and 2017 were too low and, therefore, did not yield any estimates.

APPENDIX I: Data Sources Recommended by the State Epidemiological Outcomes Work Group (SEOW)

Data Set	Source	Years	How to Access	Coverage	Target
Alcohol-Related Disease Impact (ARDI) Database	CDC	Based on averages 2006-2010	http://nccd.cdc.gov/DPH_ARDI/default/default.aspx	U.S. and states	General population
Automated Reporting Information Exchange System (ARIES)	ISP	Annual Most recent 2017	On request from ISP	Indiana and counties	Vehicle collisions in general population
Behavioral Risk Factor Surveillance System (BRFSS)	CDC	Annual Most recent 2017	http://www.cdc.gov/brfss/brfssprevalence/index.html	U.S. and states	Adults 18 and older
Hospital Discharge Database	ISDH	Annual Most recent 2017	http://www.in.gov/isdh/20624.htm	Indiana and counties	General population
Indiana Adult Tobacco Survey (IATS)	ISDH/TPCC	Bi-annual Most recent 2017		Indiana	Adults
Indiana College Substance Use Survey	ICAN/IPRC	Annual Most recent 2018	http://www.drugs.indiana.edu/indiana-college-survey/substance-use-survey	Indiana	College students
Indiana Clandestine Meth Lab Seizures	ISP	Annual Most recent 2018	On request from ISP	Indiana and counties	General population
Indiana Youth Survey	IPRC	Annual Most recent 2018	http://inys.indiana.edu/survey-results	Indiana and regions	6th – 12th grade students in Indiana
Indiana Youth Tobacco Survey (IYTS)	ISDH/TPCC	Bi-annual Most recent 2016	On request from ISDH	Indiana	6th – 12th grade students in Indiana
Monitoring the Future (MTF) Survey	NIDA	Annual Most recent 2018	http://www.monitoringthefuture.org/data/data.html	U.S.	8th, 10th, and 12th grade students
Mortality data	ISDH	Annual	On request from ISDH	Indiana and counties	General population
	CDC	Annual Most recent 2016	https://wonder.cdc.gov/	U.S., states, and counties	General population

Continued on Next Page

APPENDIX I (continued)

Data Set	Source	Years	How to Access	Coverage	Target
National Survey on Drug Use and Health (NSDUH)	SAMHSA	Annual Most recent 2017	http://www.samhsa.gov/data/population-data-nsduh	U.S., states, and some sub-state estimates	Population 12 years and older
Population Estimates	U.S. Census Bureau	Annual	http://www.census.gov/	U.S., states, and counties	General population
Treatment Episode Data Set (TEDS)	SAMHSA	Annual Most recent 2016	http://www.dasis.samhsa.gov/dasis2/teds.htm	U.S. and states; for county-level data contact Indiana DMHA	Substance abuse treatment population eligible for public services (200% FPL)
	DMHA	Annual Most recent 2018			
National Incident-Based Reporting System	FBI	Most recent 2017	https://ucr.fbi.gov/nibrs/2017/tables/data-tables	U.S., states, and counties	Arrests summarized by offense category
Uniform Reporting System (URS) – Mental Health National Outcomes Measures	SAMHSA	Annual Most recent 2016	https://www.dasis.samhsa.gov/dasis2/lurs.htm	U.S. and states	Treatment population eligible for public services (200% FPL)
			https://www.samhsa.gov/data/report/2017-uniform-reporting-system-urs-table-indiana		
Youth Risk Behavior Surveillance System (YRBSS)	CDC	Bi-annual Most recent 2017 (Indiana 2015)	http://nccd.cdc.gov/YouthOnline/App/Default.aspx	U.S. and states	High school students

Abbreviations used: CDC = Centers for Disease Control and Prevention; DMHA = Division of Mental Health & Addiction; FBI = Federal Bureau of Investigation; ICAN = Indiana Collegiate Action Network; IPRC = Indiana Prevention Resource Center; ISDH = Indiana State Department of Health; ISP = Indiana State Police; NACJD = National Archive of Criminal Justice Data; NIDA = National Institute on Drug Abuse; SAMHSA = Substance Abuse and Mental Health Services Administration; TPCC = Tobacco Prevention & Cessation Commission.

APPENDIX II: SUBSTANCE USE INDICATORS AT-A-GLANCE

SUBSTANCE	USE OR CONSEQUENCE	TARGET POPULATION	DATASET	
Alcohol	Past-month use	General population ages 12+	NSDUH	
	Past-month binge drinking			
	Alcohol use disorder in the past year	Adults ages 18+	BRFSS	
	Needing but not receiving treatment for alcohol use			
	Past-month alcohol use	Grades 9-12	YRBSS	
	Past-month binge drinking			
	Past-month heavy drinking			
	Past-month chronic drinking			
	Ever drank alcohol			
	Drank alcohol before age 13 years			
	Currently drank alcohol			
	Usually obtained the alcohol they drank by someone giving it to them			
	Drank five or more drinks of alcohol in a row			
	Reported that the largest number of drinks they had in a row was 10 or more			
Past-month alcohol use	Grades 6-12	INYS		
Past-month binge drinking				
Mean age of first use	Treatment population at or below 200% FPL, in state-sponsored programs	TEDS		
Use reported at treatment admission				
Primary use (dependence) reported at treatment admission				
Alcohol-related crashes				
Alcohol-related fatal crashes				
Alcohol-attributable deaths				
Alcohol-attributable fractions				
Years of potential life lost due to excessive alcohol use				
Alcohol-induced deaths				
Past-month use of tobacco product			General population ages 12+	NSDUH
Past-month use of Cigarettes				
Past-month smoking			Adults ages 18+	BRFSS
Past-month smokeless tobacco				
Four-level smoking status			Middle and high school students	IYTS
Past-month and lifetime use of various tobacco products				
Ever tried cigarette smoking	Grades 9-12	YRBSS		
Smoked a whole cigarette before age 13 years				
Past-month use of various tobacco products	Grades 6-12	INYS		
Past-month use of various tobacco products				
Mean age of first use	General population	IATS		
Use of various tobacco products				
Cessation intentions and attempts				
Percentage of smoke-free homes and work places				

Continued on Next Page

APPENDIX II (continued)

SUBSTANCE	USE OR CONSEQUENCE	TARGET POPULATION	DATASET		
Marijuana	Past-month use	General population ages 12+	NSDUH		
	Past-year use				
	First use				
	Ever used marijuana	Grades 9-12	YRBSS		
	Tried marijuana before age 13 years				
	Currently used marijuana				
	Ever used synthetic marijuana				
	Usually used marijuana by smoking it	Grades 6-12	INYS		
	Past-month use of marijuana and synthetic marijuana				
	Mean age of first use	Treatment population at or below 200% FPL, in state-sponsored programs	TEDS		
Use reported at treatment admission					
Primary use (dependence) reported at treatment admission					
Past-year use					
Lifetime use					
Past-month use of cocaine/crack					
Mean age of first use					
Use reported at treatment admission					
Primary use (dependence) reported at treatment admission					
Arrests for possession and sale of marijuana					
Heroin	Past-year use	General population	UCR		
	Lifetime use				
	Used a needle to inject any illegal drug at least once during their lifetime	Grades 9-12	NSDUH		
	Past-month use				
	Mean age of first use	Grades 6-12	INYS		
	Use reported at treatment admission				
	Primary use (dependence) reported at treatment admission	Treatment population at or below 200% FPL, in state-sponsored programs	TEDS		
	Past-year use				
	Lifetime use				
	Used a needle to inject any illegal drug at least once during their lifetime				
Past-month use					
Mean age of first use					
Use reported at treatment admission					
Primary use (dependence) reported at treatment admission					
Past-year use					
Lifetime use					
Methamphetamine	Past-month use	Grades 9-12	NSDUH		
	Mean age of first use				
	Use reported at treatment admission	Grades 6-12	YRBSS		
	Primary use (dependence) reported at treatment admission				
	Arrests for possession and sale of synthetic drugs	Treatment population at or below 200% FPL, in state-sponsored programs	TEDS		
	Clandestine meth lab seizures				
	Children identified/rescued in lab homes				
	Arrests made during lab seizures				
	General population			General population	UCR
	ISP Meth Lab Seizures				

Continued on Next Page

APPENDIX II (continued)

SUBSTANCE	USE OR CONSEQUENCE	TARGET POPULATION	DATASET
Prescription Drugs	Past-year misuse of pain relievers	General population ages 12+	NSDUH
	Past-month use of prescription drugs	Grades 6-12	INYS
	Mean age of first use		
	Past-year dispensation of opioids	General population	INSPECT
	Use reported at treatment admission	Treatment population at or below 200% FPL, in state-sponsored programs	TEDS
	Primary use (dependence) reported at treatment admission	General population	ISDH, CDC

Abbreviations used: ARDI = Alcohol-Related Disease Impact database; ARIES = Automated Reporting Information Exchange System; BRFSS = Behavioral Risk Factor Surveillance System; CDC = Centers for Disease Control and Prevention; IATS = Indiana Adult Tobacco Survey; INSPECT = Indiana Scheduled Prescription Drug Electronic Collection and Tracking system; INYS = Indiana Youth Survey; ISDH = Indiana State Department of Health; ISP = Indiana State Police; IYTS = Indiana Youth Tobacco Survey; NSDUH = National Survey on Drug Use and Health; SAMMEC = Smoking-Attributable Mortality, Morbidity, and Economic Costs; TEDS = Treatment Episode Data Set; UCR = Uniform Crime Reporting program; YRBSS = Youth Risk Behavior Surveillance System.

Additional information on these datasets, including how to access them, can be found in Appendix I.

APPENDIX III: CLUSTER ANALYSIS

We completed a statewide cluster analysis to determine the drug combinations that are most frequently used by polysubstance users who are in treatment. Results were based on the 2018 Treatment Episode Data Set, which we received from the Indiana Family and Social Services Administration (FSSA, 2018).

Drugs were grouped into nine (9) categories:

- Alcohol
- Marijuana
- Opioids (including nonprescription methadone, heroin, and other opiates/synthetics)
- Cocaine
- Methamphetamine
- Hallucinogens (including PCP and other hallucinogens)
- Stimulants (including amphetamines and other stimulants)
- Sedatives (including benzodiazepines, barbiturates, and sedatives/hypnotics)
- Other drugs (including inhalants, over-the-counter medications, other drugs, and unknown substances)

The analysis indicated that more than 70% of Hoosiers who received substance use treatment in 2018 reported misusing two or more drugs. Polysubstance users primarily fell into one of nine drug clusters (see Table III.1). The most commonly used combination of drugs included alcohol and marijuana. Alcohol combined with a drug that fell into the “other” drug category was the second most frequent grouping. Overall, marijuana

was the drug most commonly combined with another substance and showed up in five out of the nine drug clusters; alcohol and methamphetamine each were represented in four clusters (see Table III.1).

The demographic composition of polysubstance users differed depending on which combination of drugs they used. Males made up a greater percentage of persons in seven of the nine drug clusters; however, women were more strongly represented in the group of individuals who used (1) an opioid and methamphetamine and (2) made up nearly 50% of persons who used a combination of an opioid and a sedative.

Whites composed the majority of polysubstance users in all of the nine drug use groupings. Blacks were most strongly represented among individuals who reported using alcohol, cocaine, and marijuana, making up over one third of persons in this category. Hispanics made up less than 10% of polysubstance users across all drug combination categories.

At least half of polysubstance users in eight of the nine polysubstance groups were between the ages of 25 and 44. Polysubstance users were somewhat younger if they reported using a combination of alcohol and marijuana (see Table III.2).

Table III.1 Drug Combinations Used by Indiana Polysubstance Users (TEDS, 2018)

Drug Combinations	Number of Admissions	% of Admissions
Alcohol & Marijuana	4,148	17.6%
Alcohol & Other Drug	3,478	14.8%
Opioids & Methamphetamine	2,920	12.4%
Alcohol, Cocaine, & Marijuana	2,564	10.9%
Marijuana, Opioids, & Methamphetamine	2,528	10.7%
Opioids & Other Drug	2,452	10.4%
Marijuana & Methamphetamine	2,243	9.5%
Alcohol, Marijuana, & Methamphetamine	2,127	9.0%
Opioids & Sedatives	1,098	4.7%

Source: FSSA, 2018

Table III.2 Demographic Characteristics of Individuals within Polysubstance Groups (Treatment Episode Data Set, 2017)

	Alcohol & Marijuana		Alcohol & Other Drug		Opioids & Meth		Alcohol, Cocaine, & Marijuana		Marijuana, Opioids, & Meth	
	N	%	N	%	N	%	N	%	N	%
Gender										
Male	2,935	70.8	2,341	67.3	1,297	44.4	1,539	60.0	1,388	54.9
Female	1,213	29.2	1,137	32.7	1,623	55.6	1,025	40.0	1,140	45.1
Race										
White	3,240	78.1	2,869	82.5	2,765	94.7	1,508	58.8	2,359	93.3
Black	656	15.8	441	12.7	37	1.3	877	34.2	83	3.3
Other	252	6.1	168	4.8	118	4.0	179	7.0	86	3.4
Ethnicity										
Hispanic	291	7.0	223	6.4	116	4.0	238	9.3	112	4.4
Non-Hispanic	3,857	93.0	3,255	93.6	2,804	96.0	2,326	90.7	2,416	95.6
Age										
Under 18	199	4.8	73	2.1	5	0.2	13	0.5	22	0.9
18-24	897	21.6	515	14.8	438	15.0	269	10.5	482	19.1
25-34	1,539	37.1	1,034	29.7	1,463	50.1	819	31.9	1,335	52.8
35-44	859	20.7	822	23.6	783	26.8	642	25.0	538	21.3
45-54	402	9.7	646	18.6	190	6.5	556	21.7	125	4.9
55 and Over	252	6.1	388	11.2	41	1.4	265	10.3	26	1.0

	Opioids & Other Drug		Marijuana & Meth		Alcohol, Marijuana & Meth		Opioids & Sedatives	
	N	%	N	%	N	%	N	%
Gender								
Male	1,343	54.8	1,180	52.6	1,322	62.2	559	50.9
Female	1,109	45.2	1,063	47.4	805	37.8	539	49.1
Race								
White	2,121	86.5	2,092	93.3	1,998	93.9	994	90.5
Black	198	8.1	62	2.8	52	2.4	54	4.9
Other	133	5.4	89	4.0	77	3.6	50	4.6
Ethnicity								
Hispanic	132	4.3	90	4.6	98	4.6	57	5.2
Non-Hispanic	1,210	94.6	2,153	95.4	2,029	95.4	1,041	94.8
Age								
Under 18	49	2.0	32	1.4	19	0.9	22	2.0
18-24	465	19.0	435	19.4	248	11.7	205	18.7
25-34	1,080	44.0	931	41.5	852	40.1	501	45.6
35-44	558	22.8	572	25.5	600	28.2	254	23.1
45-54	206	8.4	227	10.1	327	15.4	70	6.4
55 and Over	94	3.8	46	2.1	81	3.8	46	4.2

Source: FSSA, 2018

MARIJUANA COCAINE PRESCRIPTION DRUGS

THE CONSUMPTION AND CONSEQUENCES OF ALCOHOL, TOBACCO, AND DRUGS IN INDIANA: A STATE EPIDEMIOLOGICAL PROFILE 2018

INDIANA STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

The Indiana State Epidemiological Outcomes Workgroup (SEOW) was established in April 2006 to review epidemiological data on the patterns and consequences of substance use and abuse in Indiana and to make recommendations to the State of Indiana regarding priorities for prevention funding for the following year. The priorities were developed based on a systematic analysis of available data, the results of which are detailed in this report.



**RICHARD M. FAIRBANKS
SCHOOL OF PUBLIC HEALTH**

INDIANA UNIVERSITY
Center for Health Policy
IUPUI



TOBACCO HEROIN METHAMPHETAMINE ALCOHOL

Our Vision

*"Healthy, safe, and drug-free environments
that nurture and assist all Indiana citizens to thrive."*

Our Mission

*"To reduce substance use and abuse
across the lifespan of Indiana citizens."*